

# UNIQUENESS AND BOUNDARY BEHAVIOUR OF LARGE SOLUTIONS TO ELLIPTIC PROBLEMS WITH SINGULAR WEIGHTS \*

M. CHUAQUI<sup>1</sup>, C. CORTÁZAR<sup>1</sup>, M. ELGUETA<sup>1</sup>  
AND J. GARCÍA-MELIÁN<sup>2,3</sup> †

<sup>1</sup> Facultad de Matemáticas, Universidad Católica de Chile,  
Casilla 306, correo 22 - Santiago, CHILE

<sup>2</sup> Dpto. de Análisis Matemático, Universidad de La Laguna,  
c/. Astrofísico Francisco Sánchez s/n, 38271 - La Laguna, SPAIN

<sup>3</sup> Centro de Modelamiento Matemático, Universidad de Chile,  
Blanco Encalada 2120, 7 piso - Santiago, CHILE

## ABSTRACT

We consider the elliptic problems  $\Delta u = a(x)u^m$ ,  $m > 1$ , and  $\Delta u = a(x)e^u$  in a smooth bounded domain  $\Omega$ , with the boundary condition  $u = +\infty$  on  $\partial\Omega$ . The weight function  $a(x)$  is assumed to be Hölder continuous, growing like a negative power of  $d(x) = \text{dist}(x, \partial\Omega)$  near  $\partial\Omega$ . We show existence and nonexistence results, uniqueness and asymptotic estimates near the boundary for both the solutions and their normal derivatives.

## 1. INTRODUCTION

Elliptic problems with blow-up on the boundary have been intensively studied in the past few years. One usually looks for a function  $u \in C^2(\Omega)$  solving an elliptic equation in a smooth bounded domain  $\Omega \subset \mathbb{R}^N$  and such that  $u(x) \rightarrow +\infty$  as  $d(x) := \text{dist}(x, \partial\Omega) \rightarrow 0+$ .

In this paper we study the problems corresponding to nonlinearities of power and of exponential type. More precisely,

$$\begin{cases} \Delta u = a(x)u^m & \text{in } \Omega \\ u = +\infty & \text{on } \partial\Omega \end{cases} \quad (P)$$

and

$$\begin{cases} \Delta v = a(x)e^v & \text{in } \Omega \\ v = +\infty & \text{on } \partial\Omega, \end{cases} \quad (P')$$

---

\*Supported by FONDECYT projects numbers 1030589 and 1030798 (Chile) and FEDER-MCYT project under contract BFM2001-3894 (Spain).

†E-mail addresses: mchuaqui@mat.puc.cl, ccortaza@mat.puc.cl, melgueta@mat.puc.cl, jjgarmel@ull.es

where  $a(x)$  is a weight function.

Problem  $(P)$  was studied for the first time in [23], with  $a \equiv 1$  and  $p = (N + 2)/(N - 2)$ . Later, in [20], [2], [3] and [27], the case with  $a(x)$  smooth and positive up to the boundary and, in [14], the case where  $a$  is smooth but vanishes on the boundary were considered. The extension to  $p$ -Laplacian equations is studied in [9]. In all these papers, uniqueness is shown by means of an estimate of the form  $u \sim Cd^{-\alpha}$  as  $d \rightarrow 0+$ . We also quote [19], where uniqueness with  $a \equiv 1$  is obtained without exact estimates for solutions on the boundary.

As for problem  $(P')$ , it was considered for the first time in [5] ( $a \equiv 1$ ), and later in [21] ( $a$  smooth and positive up to  $\partial\Omega$ ) and [1] ( $a \equiv 1$ ). In [21] and [1], uniqueness and asymptotic estimates near the boundary were also provided.

More recently, some results have appeared which treat the situation where the weight  $a$  is possibly unbounded near  $\partial\Omega$ . In [25], the radial case was considered for more general operators than the Laplacian, but obtaining only existence and nonexistence results. For problem  $(P)$  (including the range  $0 < m \leq 1$ ), the radial case was completely discussed in [6]; the authors obtained existence, nonexistence, uniqueness, multiplicity and estimates for all positive solutions. Finally, in [28] and [13], the existence of solutions in smooth bounded domains of  $\mathbb{R}^N$  was shown for both problems  $(P)$  and  $(P')$  (actually [13] admits more general power-like and exponential-like nonlinearities without monotonicity assumptions).

We mention in passing other relevant papers where related problems were considered, for instance [4], [8], [12], [18], [19], [22], [24] and [26] (see also [7], [10], [11], [15], [16] for elliptic systems).

Our goal here is to provide a complete picture of the set of solutions to problems  $(P)$  and  $(P')$  when the weight  $a(x)$  is singular on  $\partial\Omega$ . We show existence, nonexistence, uniqueness and estimates near the boundary for solutions and their normal derivatives in smooth bounded domains of  $\mathbb{R}^N$ . Techniques based on sub and supersolutions were used in [8], [14] and [16]. We will proceed in a different manner, which in particular will give us information about the growth of the normal derivatives of the solutions. For the sake of completeness we also prove existence, although by a different method than that in [28] and [13].

We will assume that  $\Omega$  is a  $C^{2,\mu}$  bounded domain for some  $0 < \mu < 1$ , and that the weight  $a(x)$  is a locally  $\mu$ -Hölder continuous function which verifies

$$C_1 d(x)^{-\gamma} \leq a(x) \leq C_2 d(x)^{-\gamma}, \quad x \in \Omega \quad (A)$$

for some positive constants  $\gamma, C_1, C_2$ . Under these assumptions, we will show that problems  $(P)$  and  $(P')$  only admit solutions when  $0 < \gamma < 2$ , in which case they are unique and satisfy bounds in terms of the data. For certain estimates near the boundary, we will sometimes need to assume the stronger condition:

$$\begin{cases} \text{there exists a bounded and positive function } C_0 \text{ defined on } \partial\Omega \\ \text{such that } \lim_{x \rightarrow x_0} d(x)^\gamma a(x) = C_0(x_0) \text{ for every } x_0 \in \partial\Omega. \end{cases} \quad (A')$$

We will see that this implies estimates of the growth of both the solutions and their normal derivatives near  $\partial\Omega$ . Note that problem  $(P)$  only makes sense for positive solutions (unless we define  $u^m$  in a suitable way for negative values of  $u$ ), while the solutions in  $(P')$  may change sign. Our results are the following.

**Theorem 1.** *Assume  $a \in C^\mu(\Omega)$  verifies hypotheses (A). Then problem (P) has no positive solutions if  $\gamma \geq 2$ , and it has a unique positive solution  $u \in C^{2,\mu}(\Omega)$  when  $0 < \gamma < 2$ . Moreover,*

$$(D_1 \sup_{\Omega} d(x)^\gamma a(x))^{-\frac{1}{m-1}} d(x)^{-\alpha} \leq u(x) \leq (D_2 \inf_{\Omega} d(x)^\gamma a(x))^{-\frac{1}{m-1}} d(x)^{-\alpha} \quad \text{in } \Omega, \quad (1)$$

where  $\alpha = (2 - \gamma)/(m - 1)$ , and  $D_1, D_2$  are positive constants depending only on  $\alpha$  and  $\Omega$ . If in addition  $a$  verifies condition (A'), then

$$\begin{aligned} \lim_{x \rightarrow x_0} d(x)^\alpha u(x) &= \left( \frac{\alpha(\alpha + 1)}{C_0(x_0)} \right)^{\frac{1}{m-1}}, \quad \lim_{x \rightarrow x_0} d(x)^{\alpha+1} \nabla u(x) \nu(x_0) = \alpha \left( \frac{\alpha(\alpha + 1)}{C_0(x_0)} \right)^{\frac{1}{m-1}}, \\ \lim_{x \rightarrow x_0} d(x)^{\alpha+2} \text{Hess } u(x) [\nu(x_0), \nu(x_0)] &= \alpha(\alpha + 1) \left( \frac{\alpha(\alpha + 1)}{C_0(x_0)} \right)^{\frac{1}{m-1}}, \end{aligned} \quad (2)$$

for every  $x_0 \in \partial\Omega$ , where  $\text{Hess } u$  denotes the Hessian of  $u$  and  $\nu$  stands for the exterior unit normal to  $\partial\Omega$ .

**Theorem 2.** *Assume  $a \in C^\mu(\Omega)$  verifies hypotheses (A). Then problem (P') has no solutions if  $\gamma \geq 2$ , and it has a unique solution  $v \in C^{2,\mu}(\Omega)$  when  $0 < \gamma < 2$ . Moreover,*

$$D'_1(2-\gamma)(\sup_{\Omega} d(x)^\gamma a(x))^{-1} d(x)^{\gamma-2} \leq e^{v(x)} \leq D'_2(2-\gamma)(\inf_{\Omega} d(x)^\gamma a(x))^{-1} d(x)^{\gamma-2} \quad \text{in } \Omega, \quad (3)$$

where  $D'_1, D'_2$  are positive constants depending only on  $\Omega$ . If in addition  $a$  verifies the asymptotic condition (A'), then

$$\begin{aligned} \lim_{x \rightarrow x_0} (v(x) + (2 - \gamma) \log d(x)) &= \log \left( \frac{2 - \gamma}{C_0(x_0)} \right), \quad \lim_{x \rightarrow x_0} d(x) \nabla v(x) \nu(x_0) = (2 - \gamma), \\ \lim_{x \rightarrow x_0} d(x)^2 \text{Hess } u(x) [\nu(x_0), \nu(x_0)] &= (2 - \gamma) \end{aligned} \quad (4)$$

for every  $x_0 \in \partial\Omega$ , where  $\nu$  stands for the exterior unit normal to  $\partial\Omega$ .

*Remarks 1.* a) Existence of solutions holds under the weaker assumption  $0 < a(x) \leq C_2 d(x)^{-\gamma}$  in  $\Omega$ , for some  $0 < \gamma < 2, C_2 > 0$ .

b) Estimate (3) implies the weaker estimate

$$\lim_{x \rightarrow x_0} \frac{v(x)}{-\log d(x)} = 2 - \gamma \quad (5)$$

for every  $x_0 \in \partial\Omega$ . Condition (A') is not needed for (5) to hold.

c) As a byproduct of the proof of estimates (2) and (4) (see Section 4) it follows that the tangential derivatives of the solutions grow at a lower rate than the normal derivatives.

d) If the weight  $a(x)$  is more regular, say  $a \in C^{k,\mu}(\Omega)$  and  $\Omega$  is  $C^{k+2,\mu}$ , for some  $k \in \mathbb{N}$ , then we can establish estimates similar to (2) and (4) for the derivatives up to order  $k + 2$  in the direction of the normal.

The paper is organized as follows: Section 2 is devoted to the existence of solutions to problems (P) and (P'). In Section 3 we obtain estimates (1) and (3) which, in particular, imply uniqueness and nonexistence when  $\gamma \geq 2$ . Finally, Section 4 deals with the proof of the estimates (2) and (4). Theorems 1 and 2 will then follow.

## 2. EXISTENCE OF SOLUTIONS

Existence of solutions of problems  $(P)$  and  $(P')$  is already known from [28] and [13] when  $0 < \gamma < 2$ . We give here a different proof based on a standard technique. We construct solutions with finite datum  $n \in \mathbb{N}$  and then obtain the desired solution by passing to the limit as  $n \rightarrow +\infty$ . The key step in this process is to show local uniform bounds.

**Lemma 3.** *Let  $n \in \mathbb{N}$ , and assume  $a \in C^\mu(\Omega)$  verifies (A) with  $0 < \gamma < 2$ . Then the problem*

$$\begin{cases} \Delta u = a(x)u^m & \text{in } \Omega \\ u = n & \text{on } \partial\Omega \end{cases} \quad (P_n)$$

*admits a unique positive solution  $u_n \in C^{2,\mu}(\Omega) \cap C(\bar{\Omega})$ . Moreover,  $u_n$  is increasing in  $n$ .*

*Proof.* Choose a cut-off function  $\varphi \in C^1(\mathbb{R}^+)$  such that  $0 \leq \varphi \leq 1$ ,  $\varphi(t) = 0$  if  $0 \leq t \leq 1$ ,  $\varphi(t) = 1$  for  $t \geq 2$ , and for  $k \in \mathbb{N}$  set  $a_k(x) = a(x)\varphi(kd(x))$ . Then  $a_k \in C^\mu(\bar{\Omega})$  (we recall that  $d(x)$  is always Lipschitz-continuous). Consider the problem

$$\begin{cases} \Delta w = a_k(x)(w+n)^m & \text{in } \Omega \\ w = 0 & \text{on } \partial\Omega. \end{cases} \quad (6)$$

Since we can take  $\underline{w} = -n$ ,  $\bar{w} = 0$  as ordered sub and supersolutions, respectively, the existence of at least a solution  $w_k^n \in C^{2,\mu}(\bar{\Omega})$  is established. By the monotonicity of the right-hand side and the maximum principle it follows that  $w_k^n$  is unique.

In order to pass to the limit as  $k \rightarrow +\infty$  in (6) we need the following lemma which appears in [17] (see Lemma 4.9 and Problem 4.6).

**Lemma 4.** *Let  $\Omega$  be a  $C^2$  bounded domain of  $\mathbb{R}^N$  and  $f \in C^\mu(\Omega)$  such that  $\sup_\Omega d(x)^\gamma |f(x)| < +\infty$  for some  $1 < \gamma < 2$ . Then the problem  $\Delta u = f$  in  $\Omega$  with  $u = 0$  on  $\partial\Omega$  has a unique solution  $u \in C^{2,\mu}(\Omega) \cap C(\bar{\Omega})$ , and*

$$\sup_\Omega d(x)^{\gamma-2} |u(x)| \leq C \sup_\Omega d(x)^\gamma |f(x)|,$$

*where  $C$  is a positive constant depending only on  $\Omega$  and  $\gamma$ .*

*Remark 2.* Notice that if  $\sup_\Omega d(x)^\gamma |f(x)| < +\infty$ , we also have  $\sup_\Omega d(x)^{\gamma'} |f(x)| < +\infty$  for every  $\gamma' > \gamma$ . In particular, when  $0 < \gamma \leq 1$ , Lemma 4 is still applicable by choosing some  $\gamma'$  such that  $1 < \gamma' < 2$ , but the estimate we obtain now is  $|u(x)| \leq Cd(x)^{2-\gamma'}$ ,  $x \in \Omega$ , which is not the optimal one.

We now apply this lemma to the problem (6). Since  $0 \leq a_k(x) \leq a(x) \leq C_2 d(x)^{-\gamma}$ , and  $w_k^n$  are uniformly bounded independently of  $k$ , we have

$$\sup_\Omega d(x)^{\gamma-2} |w_k^n(x)| \leq C \quad (7)$$

for some constant  $C$  not depending on  $k$  (this holds true in the whole range  $0 < \gamma < 2$ , cf. Remark 2). The uniform bounds for the sequence  $\{w_k^n\}$  and a standard bootstrapping argument give local  $C^{2,\mu}$  bounds, so that by means of a diagonal procedure we can obtain

a convergent subsequence  $w_k^n \rightarrow w_n$  in  $C_{\text{loc}}^2(\Omega)$  as  $k \rightarrow +\infty$ . Passing to the limit in (6) we obtain that  $\Delta w_n = a(x)(w_n + n)^m$  in  $\Omega$ , and  $\sup_{\Omega} d(x)^{\gamma-2}|w_n(x)| \leq C$  thanks to (7). Since  $0 < \gamma < 2$ , this implies  $w_n \in C(\bar{\Omega})$  and  $w_n = 0$  on  $\partial\Omega$ . The function  $u_n = w_n + n$  is then the required solution to  $(P_n)$  (notice that  $u_n$  is positive, since  $-n < w_n < 0$ ). By elliptic regularity,  $u_n \in C^{2,\mu}(\Omega)$ . Uniqueness and monotonicity with respect to  $n$  are a consequence of the maximum principle.  $\square$

The following lemma is the equivalent to Lemma 3 for  $f(u) = e^u$ . We omit the proof, which is essentially the same. The only point worth stressing is that we can find a subsolution to the truncated problem of the form  $\underline{u} = A(|x|^2 - B)$  for sufficiently large  $A$  and  $B$ , and that uniform bounds for  $e^{w_k^n}$  are enough to pass to the limit via Lemma 4.

**Lemma 5.** *Let  $n \in \mathbb{N}$ , and assume  $a \in C^\mu(\Omega)$  verifies (A) with  $0 < \gamma < 2$ . Then the problem*

$$\begin{cases} \Delta v = a(x)e^v & \text{in } \Omega \\ v = n & \text{on } \partial\Omega \end{cases} \quad (P'_n)$$

*admits a unique solution  $v_n \in C^{2,\mu}(\Omega) \cap C(\bar{\Omega})$ . Moreover,  $v_n$  is increasing in  $n$ .*

We finally prove that the sequences  $\{u_n\}$  and  $\{v_n\}$  constructed above converge, as  $n \rightarrow +\infty$ , to solutions of problems  $(P)$  and  $(P')$ , respectively.

**Lemma 6.** *Let  $\{u_n\}$ ,  $\{v_n\}$  be the sequences of solutions to problems  $(P_n)$  and  $(P'_n)$  given by Lemma 3 and Lemma 5, respectively. Then  $u_n \rightarrow u$  in  $C_{\text{loc}}^2(\Omega)$ , where  $u \in C^{2,\mu}(\Omega)$  is a positive solution to  $(P)$  and  $v_n \rightarrow v$  in  $C_{\text{loc}}^2(\Omega)$ , where  $v \in C^{2,\mu}(\Omega)$  is a solution to  $(P')$ .*

*Proof.* Since  $a$  is strictly positive in  $\Omega$ , there exists  $a_0 > 0$  such that  $a \geq a_0$ . Then  $u_n$  verifies  $\Delta u_n \geq a_0 u_n^m$  in  $\Omega$ , and the maximum principle implies  $u_n \leq U$ , where  $U$  is the unique solution to problem  $(P)$  with  $a(x) \equiv a_0$  (cf. [3]). This gives local uniform bounds for  $u_n$ , and an argument similar to the used in the proof of Lemma 3 gives that  $u_n \rightarrow u$  in  $C_{\text{loc}}^2(\Omega)$  for some function  $u$ . Passing to the limit in  $(P_n)$  we see that  $\Delta u = a(x)u^m$ , and since  $u_n$  is increasing it also follows that  $u = +\infty$  on  $\partial\Omega$ . Finally, by elliptic regularity,  $u \in C_{\text{loc}}^{2,\mu}(\Omega)$ , and is a classical solution to  $(P)$ . The proof for the sequence  $\{v_n\}$  is similar.  $\square$

### 3. GLOBAL ESTIMATES AND UNIQUENESS

This section is devoted to prove the global estimates (1) and (3), the nonexistence result for  $\gamma \geq 2$  and the uniqueness of solutions to problems  $(P)$  and  $(P')$ .

**Theorem 7.** *Assume  $a \in C^\mu(\Omega)$  verifies hypothesis (A). Then there exist positive constants  $D_1, D_2$  depending only on  $\alpha$  and  $\Omega$ , such that*

$$(D_1 \sup_{\Omega} d(x)^\gamma a(x))^{-\frac{1}{m-1}} d(x)^{-\alpha} \leq u(x) \leq (D_2 \inf_{\Omega} d(x)^\gamma a(x))^{-\frac{1}{m-1}} d(x)^{-\alpha} \quad \text{in } \Omega, \quad (1)$$

*for every positive solution  $u$  to  $(P)$ .*

*Proof.* Let  $u$  be a positive solution to (P). We show first that  $\inf_{\Omega} d(x)^{\alpha} u(x) > 0$ ,  $\sup_{\Omega} d(x)^{\alpha} u(x) < +\infty$ .

Consider an arbitrary sequence  $\{x_n\} \subset \Omega$  such that  $d_n := d(x_n) \rightarrow 0$ . In the ball of center  $x_n$  and radius  $d_n/2$  we have  $\Delta u \geq C_1(3/2)^{-\gamma} d_n^{-\gamma} u^m$ . Setting

$$u_n(x) = \left( \frac{2^{\gamma-2} C_1}{3^{\gamma}} \right)^{\frac{1}{m-1}} d_n^{\alpha} u(x_n + \frac{d_n}{2} x), \quad x \in B(0, 1),$$

we obtain  $\Delta u_n \geq u_n^m$  in  $B(0, 1)$ . By the maximum principle,  $u_n(x) \leq U(x)$ ,  $x \in B(0, 1)$ , where  $U$  stands for the unique positive solution to  $\Delta U = U^m$  in  $B(0, 1)$ ,  $U|_{\partial B(0,1)} = +\infty$ . For  $x = 0$  we obtain

$$d_n^{\alpha} u(x_n) \leq \left( \frac{3^{\gamma}}{2^{\gamma-2} C_1} \right)^{\frac{1}{m-1}} U(0).$$

Since the sequence  $\{x_n\}$  is arbitrary, this shows that  $\sup_{\Omega} d(x)^{\alpha} u(x) < +\infty$ .

To prove the lower bound, we let  $v = u^{-p}$  with  $0 < p < m - 1$  to be chosen. Then

$$-\Delta v + \frac{p+1}{p} \frac{|\nabla v|^2}{v} = p a(x) v^{-r},$$

where  $r = (m - 1 - p)/p > 0$ . It follows that  $-\Delta v \leq C d(x)^{-2+p\alpha}$  in  $\Omega$  ( $C$  will always denote a positive constant). We now choose  $p$  small enough so that  $1 < 2 - p\alpha < 2$ . Let  $\phi$  be the unique solution to the problem  $-\Delta \phi = C d(x)^{-2+p\alpha}$ ,  $\phi|_{\partial\Omega} = 0$ , which exists due to Lemma 4. By the maximum principle and Lemma 4,  $v \leq \phi \leq C d(x)^{p\alpha}$ , that is  $u(x) \geq C d(x)^{-\alpha}$  in  $\Omega$ .

We finally prove (1). Set  $p = 1 + 2/\alpha$ , and let  $\lambda > 0$  to be determined later. Since  $m - p < 0$ , we have

$$\begin{aligned} \Delta(\lambda u) &= \lambda^{1-p} d(x)^{\gamma} a(x) (d(x)^{\alpha} u(x))^{m-p} (\lambda u)^p \\ &\geq \lambda^{1-p} \inf_{\Omega} d(x)^{\gamma} a(x) (\sup_{\Omega} d(x)^{\alpha} u(x))^{m-p} (\lambda u)^p \\ &= (\lambda u)^p \quad \text{in } \Omega, \end{aligned}$$

if  $\lambda = ((\inf_{\Omega} d(x)^{\gamma} a(x)) (\sup_{\Omega} d(x)^{\alpha} u(x))^{m-p})^{\frac{1}{p-1}}$  (compare with [13]). Hence  $\lambda u$  is a subsolution to the problem  $\Delta U = U^p$  in  $\Omega$ ,  $U|_{\partial\Omega} = +\infty$ , which has a unique solution  $U$ . Since  $MU$  is a supersolution and  $\lambda u \leq MU$  in  $\Omega$  for large  $M$  (observe that  $U \sim C d^{-\alpha}$  near  $\partial\Omega$ , so that the inequality can be achieved), the method of sub and supersolutions (cf. Lemma 4 in [14] and Lemma 2 in [13]) yields  $\lambda u \leq U$ . Note that the maximum principle could not be used directly, since we do not know a priori the relation between  $\lambda u$  and  $U$  near the boundary. A little algebra now shows that this leads indeed to the upper estimate in (1). The lower estimate is proved in a similar way.  $\square$

The following theorem is the analogue of Theorem 7 in the exponential case. The proof is essentially the same and will be omitted.

**Theorem 8.** *Assume  $a \in C^{\mu}(\Omega)$  verifies hypothesis (A). Then there exist positive constants  $D'_1, D'_2$  depending only on  $\alpha$  and  $\Omega$ , such that*

$$D'_1(2-\gamma)(\sup_{\Omega} d(x)^{\gamma} a(x))^{-1} d(x)^{\gamma-2} \leq e^{v(x)} \leq D'_2(2-\gamma)(\inf_{\Omega} d(x)^{\gamma} a(x))^{-1} d(x)^{\gamma-2} \quad \text{in } \Omega, \quad (3)$$

for every solution  $v$  to (P').

For  $\gamma \geq 2$  the estimates (1) and (3) imply that all solutions are bounded in  $\bar{\Omega}$ . We thus have the following corollary. See Lemma 5.1 in [16] for a related result.

**Corollary 9.** *Assume  $a \in C^\mu(\Omega)$  verifies hypothesis (A) and  $\gamma \geq 2$ . Then problems (P) and (P') have no solutions.*

We now study the uniqueness of solutions of problems (P) and (P'). Observe that (1) and (3) are not enough to conclude that the quotients of two positive solutions to (P) tend to 1 at  $\partial\Omega$ , which has been often used to prove uniqueness. Instead we adapt the argument in Theorem 3.4 of [19].

**Theorem 10.** *Assume  $a \in C^\mu(\Omega)$  verifies (A) and  $0 < \gamma < 2$ . Then problem (P) has a unique positive solution.*

*Proof.* Let  $u, v$  be positive solutions to (P), and assume there exist  $x_0 \in \Omega$  and  $k > 1$  such that  $u(x_0) > kv(x_0)$ . Let  $\Omega_0 := \{u > kv\} \cap B_r(x_0)$ , where  $r = d(x_0)/2$ . By (1) we have

$$\Delta(u - kv) > a(x)(k^{m-1} - 1)kv^m \geq \kappa_1 r^{-\alpha-2}k$$

in  $\Omega_0$ , where  $\kappa_1$  is some positive constant. Set  $w(x) = (\kappa_1 r^{-\alpha-2}k)(r^2 - |x - x_0|^2)/2N$ . Then  $\Delta(u - kv + w) \geq 0$  in  $\Omega_0$ , and from the maximum principle we deduce the existence of  $x_1 \in \partial\Omega_0$  such that

$$u(x_0) - kv(x_0) + w(x_0) \leq u(x_1) - kv(x_1) + w(x_1).$$

It is easy to see that  $x_1 \in \partial B_r(x_0)$ . Then we obtain that  $(\kappa_1 r^{-\alpha}k)/2N \leq u(x_1) - kv(x_1)$ , which together with the upper estimate for  $v$ , leads to the existence of  $\kappa_2 > 0$  such that  $u(x_1) > (1 + \kappa_2)kv(x_1)$ . Proceeding inductively, we find a sequence  $\{x_n\} \subset \Omega$  such that  $u(x_n) > (1 + \kappa_2)^n kv(x_n)$ , which contradicts the fact that the quotients of any two positive solutions must be bounded. Hence,  $u \leq v$ , and interchanging the roles of  $u$  and  $v$  we obtain  $u \equiv v$ . This proves the theorem.  $\square$

**Theorem 11.** *Assume  $a \in C^\mu(\Omega)$  verifies (A) and  $0 < \gamma < 2$ . Then problem (P') has a unique solution.*

*Proof.* The argument is similar to the one used in Theorem 10. Let  $u, v$  be solutions to (P') and notice that (3) implies that  $u - v$  is bounded. Assume there exist  $x_0 \in \Omega$  and  $k > 0$  such that  $u > v + k$ . Define  $\Omega_0 := \{u > v + k\} \cap B_r(x_0)$ , for  $r = d(x_0)/2$ . Then

$$\Delta(u - v - k) > a(x)e^v(e^k - 1) \geq \kappa_1 r^{-2}$$

in  $\Omega_0$ . Setting  $w(x) = (\kappa_1 r^{-2})(r^2 - |x - x_0|^2)/2N$ , we obtain as before the existence of  $x_1 \in \partial B_r(x_0)$  such that  $w(x_0) < u(x_1) - v(x_1) - k$ . In particular,  $u(x_1) > v(x_1) + k + \kappa_1/2N$ . Iterating this procedure, we obtain a sequence  $\{x_n\} \subset \Omega$  such that  $u(x_n) > v(x_n) + k + n\kappa_2/2N$ , which contradicts the boundedness of  $u - v$ . Thus,  $u \leq v$  and similarly  $u \geq v$ , which proves uniqueness.  $\square$

*Remarks 3.* a) For problem  $(P')$ , we can use a standard procedure to show uniqueness, since the global estimate (3) shows that the quotient of two arbitrary solutions tends to one as we approach the boundary of  $\Omega$  (cf. Remarks 1 b)). Indeed, let  $u, v$  be solutions to  $(P')$ . By replacing  $u, v$  by  $u + 1, v + 1$  and  $a(x)$  by  $a(x)e^{-1}$ , we can assume that  $u, v \geq 1$ . Let  $w = u/v$ . Then

$$v\Delta w + 2\nabla v \nabla w = a(x)e^v(e^{(w-1)v} - w) \quad \text{in } \Omega.$$

Set  $\Omega^+ := \{x \in \Omega : w(x) > 1\}$ , and assume  $\Omega^+ \neq \emptyset$  (observe that  $w = 1$  on  $\partial\Omega$ , and this implies  $w = 1$  on  $\partial\Omega^+$ ). Then since the function  $h(\tau) = e^{(\tau-1)v} - \tau$  is positive for  $\tau \geq 1$ , we have  $v\Delta w + 2\nabla v \nabla w \geq 0$  in  $\Omega^+$ , and by the maximum principle  $w \leq 1$  in  $\Omega^+$ , a contradiction. Thus  $\Omega^+ = \emptyset$ , i.e.  $w \leq 1$  in  $\Omega$ , and the symmetric argument gives  $w \equiv 1$ , which shows uniqueness.

b) The proof of Theorems 10 and 11 is also valid for *unbounded* domains, as long as we have estimates like (1) and (3) for all solutions (this will be used in Section 4).

#### 4. ESTIMATES NEAR THE BOUNDARY

In this final section, we are obtaining estimates near the boundary both for solutions to  $(P)$  and  $(P')$  and their normal derivatives. Our proof is based on a rescaling argument. We will assume that  $a \in C^\mu(\Omega)$  verifies hypotheses  $(A)$  for  $0 < \gamma < 2$  and  $(A')$ .

We are performing all the calculations for problem  $(P)$ , since the translation of the argument to problem  $(P')$  is straightforward.

*Proof of estimates (2).* Let  $x_0 \in \partial\Omega$ , and  $\{x_n\} \subset \Omega$  such that  $x_n \rightarrow x_0$ . Choose an open neighbourhood  $\mathcal{U}$  of  $x_0$  so that  $\partial\Omega$  admits  $C^{2,\mu}$  local coordinates  $\xi : \mathcal{U} \rightarrow \mathbb{R}^N$ , and  $x \in \mathcal{U} \cap \Omega$  if and only if  $\xi_1(x) > 0$  ( $\xi = (\xi_1, \xi_2, \dots, \xi_N)$ ). We can moreover assume  $\xi(x_0) = 0$ . If  $u(x) = \bar{u}(\xi(x))$ , then we have the equation

$$\sum_{i,j=1}^N a_{ij}(\xi) \frac{\partial^2 \bar{u}}{\partial \xi_i \partial \xi_j} + \sum_{i=1}^N b_i(\xi) \frac{\partial \bar{u}}{\partial \xi_i} = \bar{a}(\xi) \bar{u}^m$$

in  $\xi(\mathcal{U} \cap \Omega)$ , where  $a(x) = \bar{a}(\xi(x))$ ,  $a_{ij}, b_i$  are (at least)  $C^\mu$ , and  $a_{ij}(0) = \delta_{ij}$ . Denote by  $\eta_n$  the projections onto  $\xi(\mathcal{U} \cap \partial\Omega)$  of  $\xi(x_n)$ , and introduce the functions

$$u_n(y) = d_n^\alpha \bar{u}(\eta_n + d_n y),$$

where  $d_n = d(\xi(x_n))$ . Notice that  $\xi(x_n) = \eta_n + d_n(1, 0, \dots, 0)$ . The function  $u_n$  satisfies the equation

$$\sum_{i,j=1}^N a_{ij}(\eta_n + d_n y) \frac{\partial u_n}{\partial \xi_i \partial \xi_j} + d_n \sum_{i=1}^N b_i(\eta_n + d_n y) \frac{\partial u_n}{\partial \xi_i} = d_n^\gamma \bar{a}(\eta_n + d_n y) u_n^m.$$

On the other hand, estimates (1) imply that, for  $y$  in compact subsets  $K$  of  $D := \{y \in \mathbb{R}^N : y_1 > 0\}$ , there exists  $n_0 = n_0(K)$  such that  $Cy_1^{-\alpha} \leq u_n(y) \leq C'y_1^{-\alpha}$  for  $n \geq n_0$ , where  $C, C'$  are positive constants not depending on  $K$ . These estimates, together with the

equation, a bootstrap argument and a diagonal procedure, allow us to obtain a subsequence (still labelled by  $\{u_n\}$ ) such that  $u_n \rightarrow u_0$  in  $C_{\text{loc}}^2(D)$ . In particular, we obtain that

$$\begin{cases} \Delta u_0 = C_0(x_0)y_1^{-\gamma}u_0^m & \text{in } D \\ C y_1^{-\alpha} \leq u_0 \leq C' y_1^{-\alpha} . \end{cases}$$

Thanks to Theorem 10 and Remarks 3 b), we see that this problem has a unique positive solution, which can be checked to be

$$u_0(y) = \left( \frac{\alpha(\alpha+1)}{C_0(x_0)} \right)^{\frac{1}{m-1}} y_1^{-\alpha} .$$

Thus, taking  $y = (1, 0, \dots, 0)$ , we arrive at

$$\begin{aligned} d_n^\alpha u(x_n) &\rightarrow \left( \frac{\alpha(\alpha+1)}{C_0(x_0)} \right)^{\frac{1}{m-1}} , & d_n^{\alpha+1} \frac{\partial u}{\partial \xi_1}(x_n) &\rightarrow -\alpha \left( \frac{\alpha(\alpha+1)}{C_0(x_0)} \right)^{\frac{1}{m-1}} , \\ d_n^{\alpha+2} \frac{\partial^2 u}{\partial \xi_1^2}(x_n) &\rightarrow \alpha(\alpha+1) \left( \frac{\alpha(\alpha+1)}{C_0(x_0)} \right)^{\frac{1}{m-1}} . \end{aligned}$$

This proves (2), since the sequence  $\{x_n\}$  is arbitrary.

Finally notice that if  $a(x) \in C^{k,\mu}(\Omega)$  and  $\Omega$  is  $C^{k+2,\mu}$ , for some  $k \in \mathbb{N}$  (Remarks 1 d)), we obtain  $u_n \rightarrow u_0$  in  $C_{\text{loc}}^{k+2}(\Omega)$ , so we can establish estimates for the derivatives up to order  $k+2$ .  $\square$

*Remark 4.* In case the function  $C_0(x_0)$  is constant on  $\partial\Omega$ , there is another way of producing the first estimate in (2) for problem (P). Indeed, setting  $\Omega_\delta := \{x \in \Omega : d(x) < \delta\}$  for positive small  $\delta$  and  $p = 1 + 2/\alpha$ , we have

$$\Delta(\lambda u) \leq \lambda^{1-p} \sup_{\Omega_\delta} (d(x)^\gamma a(x)) (\inf_{\Omega_\delta} d(x)^\alpha u(x))^{m-p} (\lambda u)^p$$

in  $\Omega_\delta$ . Choosing  $\lambda = \lambda(\delta) := (\sup_{\Omega_\delta} d(x)^\gamma a(x) (\inf_{\Omega_\delta} d(x)^\alpha u(x))^{m-p})^{\frac{1}{p-1}}$ , we obtain  $\Delta(\lambda u) \leq (\lambda u)^p$  in  $\Omega_\delta$ . A similar calculation as in [8] shows then that for a suitable constant  $C > 0$ ,  $\lambda u + C\delta^{-\alpha} \geq U$  in  $\Omega_\delta$ , where  $U$  stands for the unique positive solution to  $\Delta U = U^p$  in  $\Omega$ ,  $U = +\infty$  on  $\partial\Omega$ . After some manipulations, this leads to

$$\liminf_{d \rightarrow 0} d(x)^\alpha u(x) \geq \left( \frac{\alpha(\alpha+1)}{C_0} \right)^{\frac{1}{m-1}} ,$$

and a similar reasoning proves the other inequality.

As can be checked, this proof can not be adapted to nonconstant  $C_0$ , since it is global in nature, and does not provide any information on the growth of the normal derivatives.

## REFERENCES

- [1] C. BANDLE, M. ESSÈN, *On the solutions of quasilinear elliptic problems with boundary blow-up*, Sympos. Math. **35** (1994), 93–111.
- [2] C. BANDLE, M. MARCUS, *Sur les solutions maximales de problèmes elliptiques non linéaires: bornes isopérimétriques et comportement asymptotique*, C. R. Acad. Sci. Paris Sér. I Math. **311** (1990), 91–93.
- [3] C. BANDLE, M. MARCUS, *'Large' solutions of semilinear elliptic equations: Existence, uniqueness and asymptotic behaviour*, J. Anal. Math. **58** (1992), 9–24.
- [4] C. BANDLE, M. MARCUS, *On second order effects in the boundary behaviour of large solutions of semilinear elliptic problems*, Differential Integral Equations **11** (1) (1998), 23–34.
- [5] L. BIEBERBACH,  *$\Delta u = e^u$  und die automorphen Funktionen*, Math. Ann. **77** (1916), 173–212.
- [6] M. CHUAQUI, C. CORTÁZAR, M. ELGUETA, C. FLORES, J. GARCÍA-MELIÁN, R. LETELIER, *On an elliptic problem with boundary blow-up and a singular weight: the radial case*, submitted for publication.
- [7] N. DANCER, Y. DU, *Effects of certain degeneracies in the predator-prey model*, SIAM J. Math. Anal. **34** (2) (2002), 292–314.
- [8] M. DEL PINO, R. LETELIER, *The influence of domain geometry in boundary blow-up elliptic problems*, Nonlinear Anal. **48** (6) (2002), 897–904.
- [9] G. DÍAZ, R. LETELIER, *Explosive solutions of quasilinear elliptic equations: Existence and uniqueness*, Nonlinear Anal. **20** (1993), 97–125.
- [10] Y. DU, *Effects of a degeneracy in the competition model. Part I: classical and generalized steady-state solutions*, J. Diff. Eqns. **181** (2002), 92–132.
- [11] Y. DU, *Effects of a degeneracy in the competition model. Part II: perturbation and dynamical behaviour*, J. Diff. Eqns. **181** (2002), 133–164.
- [12] Y. DU, Q. HUANG, *Blow-up solutions for a class of semilinear elliptic and parabolic equations*, SIAM J. Math. Anal. **31** (1999), 1–18.
- [13] J. GARCÍA-MELIÁN, *A remark on the existence of positive large solutions via sub and supersolutions*, submitted for publication.
- [14] J. GARCÍA-MELIÁN, R. LETELIER-ALBORNOZ, J. SABINA DE LIS, *Uniqueness and asymptotic behaviour for solutions of semilinear problems with boundary blow-up*, Proc. Amer. Math. Soc. **129** (2001), no. 12, 3593–3602.
- [15] J. GARCÍA-MELIÁN, R. LETELIER-ALBORNOZ, J. SABINA DE LIS, *The solvability of an elliptic system under a singular boundary condition*, preprint.

- [16] J. GARCÍA-MELIÁN, A. SUÁREZ, *Existence and uniqueness of positive large solutions to some cooperative elliptic systems*, *Advanced Nonlinear Studies* **3** (2003), 75–88.
- [17] D. GILBARG, N.S. TRUDINGER, *Elliptic partial differential equations of second order*, Springer–Verlag, 1983.
- [18] J. B. KELLER, *On solutions of  $\Delta u = f(u)$* , *Comm. Pure Appl. Math.* **10** (1957), 503–510.
- [19] S. KIM, *A note on boundary blow-up problem of  $\Delta u = u^p$* , IMA preprint No. 1820, (2002).
- [20] V. A. KONDRAT'EV, V. A. NIKISHKIN, *Asymptotics, near the boundary, of a solution of a singular boundary value problem for a semilinear elliptic equation*, *Differential Equations* **26** (1990), 345–348.
- [21] A. C. LAZER, P. J. MCKENNA, *On a problem of Bieberbach and Rademacher*, *Nonlinear Anal.* **21** (1993), 327–335.
- [22] A. C. LAZER, P. J. MCKENNA, *Asymptotic behaviour of solutions of boundary blow-up problems*, *Differential Integral Equations* **7** (1994), 1001–1019.
- [23] C. LOEWNER, L. NIRENBERG, *Partial differential equations invariant under conformal of projective transformations*, *Contributions to Analysis* (a collection of papers dedicated to Lipman Bers), Academic Press, New York, 1974, p. 245–272.
- [24] M. MARCUS, L. VÉRON, *Uniqueness and asymptotic behaviour of solutions with boundary blow-up for a class of nonlinear elliptic equations*, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **14** (2) (1997), 237–274.
- [25] M. MOHAMMED, G. PORCU, G. PORRU, *Large solutions to some non-linear O.D.E. with singular coefficients*, *Nonlinear Anal.* **47** (2001), 513–524.
- [26] R. OSSERMAN, *On the inequality  $\Delta u \geq f(u)$* , *Pacific J. Math.* **7** (1957), 1641–1647.
- [27] L. VÉRON, *Semilinear elliptic equations with uniform blowup on the boundary*, *J. Anal. Math.* **59** (1992), 231–250.
- [28] Z. ZHANG, *A remark on the existence of explosive solutions for a class of semilinear elliptic equations*, *Nonlinear Anal.* **41** (2000), 143–148.