

# KELLER-OSSERMAN TYPE CONDITIONS FOR SOME ELLIPTIC PROBLEMS WITH GRADIENT TERMS

SALOMÓN ALARCÓN, JORGE GARCÍA-MELIÁN AND ALEXANDER QUAAS

ABSTRACT. In this paper we consider the elliptic boundary blow-up problems

$$\begin{cases} \Delta u \pm g(|\nabla u|) = f(u) & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega$  is a smooth bounded domain and the functions  $f$  and  $g$  are increasing and continuous. Our main concern will be to prove both existence and nonexistence of nonnegative solutions, depending on new integral conditions of Keller-Osserman type involving  $f$  and  $g$ . We show in particular that the problem with a minus sign may have solutions inclusive for some functions  $g$  with slightly superquadratic growth at infinity that is somehow not expected. We also obtain uniqueness of nonnegative solutions in some cases.

## 1. INTRODUCTION

The main objective of this paper is to analyze the existence and nonexistence of nonnegative solutions to the problems

$$(P_{\pm}) \quad \begin{cases} \Delta u \pm g(|\nabla u|) = f(u) & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega$  is a  $C^2$  bounded domain of  $\mathbb{R}^N$  and the functions  $f, g$  are continuous and increasing, with  $f(0) = g(0) = 0$ . By a solution to  $(P_{\pm})$  we mean a function  $u \in C^1(\Omega)$  which verifies the equation in the weak sense and  $u(x) \rightarrow \infty$  as  $d(x) := \text{dist}(x, \partial\Omega) \rightarrow 0$ .

The model problem without the gradient terms, namely

$$(1.1) \quad \begin{cases} \Delta u = f(u) & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

has generated a good deal of research. The important case  $f(u) = e^u$  was analyzed in the early works of Bieberbach [9] when  $N = 2$  and Rademacher [49] for  $N = 3$ , and later reconsidered in [36]. The other significant example  $f(u) = u^p$ ,  $p > 1$ , was dealt with in [47], [41] and [32] (also in [17] when the Laplacian is replaced by the  $p$ -Laplacian  $\Delta_p$ ). These cases have been also studied when the underlying domain  $\Omega$  is not necessarily smooth: see [44], [45] and [18].

As for general increasing nonlinearities  $f(u)$ , it has been known since the pioneering works of Keller [31] and Osserman [46] that problem (1.1) admits a solution if and only if the nowadays called Keller-Osserman condition holds, i. e.

$$(1.2) \quad \int_1^{\infty} \frac{ds}{\sqrt{F(s)}} < \infty,$$

where  $F(s) = \int_0^s f(t)dt$  (see also [20] when  $f$  is not increasing). Later, other important questions concerning boundary behavior of solutions and uniqueness or multiplicity, have been considered in many works. We quote for instance [5], [37] and [25] for uniqueness for general nonlinearities and [1], [2] for multiplicity results. See also [7], [8], [37], [15] and [3],

where more precise information on the asymptotic behavior of the solutions is obtained. A great amount of works have been also interested in the appearance of weights in the equation, both vanishing on  $\partial\Omega$  ([19], [26], [13], [14], [12], [42], [43], [24]) or singular at  $\partial\Omega$  ([10], [11], [23]). More references can be found in the survey [50].

With regard to boundary blow-up problems containing gradient terms, Lasry and Lions [35] considered the following:

$$(1.3) \quad \begin{cases} \Delta u - |\nabla u|^p = \lambda u + h & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

where  $p > 1$ ,  $\lambda > 0$  and  $h$  is smooth in  $\Omega$  (it appears in stochastic control problems with state constraints). Assuming  $h$  has a prescribed growth near  $\partial\Omega$ , it was shown in [35] that there exists a unique solution to (1.3) when  $1 < p \leq 2$ . Further developments of this and related problems have been made in [48], [38], [39]. In particular, in this paper we can extend the results in [35] when we consider  $h = 0$  and we replace the term  $|\nabla u|^p$  by  $g(|\nabla u|) = |\nabla u|^2 \log(|\nabla u| + 1)$  which is slightly superquadratic. We believe that this can be extended to cover the case of nontrivial  $h$ .

Another type of problems including gradient terms in the equation was proposed by Bandle and Giarrusso in [4]. They considered

$$(1.4) \quad \begin{cases} \Delta u \pm |\nabla u|^q = f(u) & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

for  $q > 0$  and general differentiable functions  $f$ , paying special attention to the two ‘‘classical’’ nonlinearities  $f(u) = u^p$ ,  $p > 1$  and  $f(u) = e^u$ . Some existence results were obtained, together with boundary behavior of positive solutions in most cases. This problem was later analyzed again in [28] and [29] (see also some extensions to problems containing weights in [27], [34], [52], [53] and [54]).

However, at the best of our knowledge, no results are available for the general problems  $(P_{\pm})$  aside the case  $g(t) = t^q$ ,  $q > 0$ . Thus in the present paper, and under quite general assumptions on  $f$  and  $g$ , we will obtain conditions to ensure existence or nonexistence of nonnegative solutions. Let us mention that our conditions are not sharp, due to the fact that the one dimensional version of  $(P_{\pm})$  is not integrable, but nevertheless they are in the case of power nonlinearities  $f(t) = t^p$ ,  $g(t) = t^q$ ,  $p, q > 0$ .

We would like to stress that for problem (1.4) with the minus sign, the results in [4] show that the exponent  $q = 2$  is critical in some sense, since no solutions are expected to exist when  $q > 2$  (although this fact is not proved there for smooth bounded domains). Among other things, we show that for problem  $(P_-)$  it is possible to have solutions for nonlinearities which grow faster than quadratically, the essential feature being an integrability condition on  $s/g(s)$  at infinity.

In our results about existence we need to obtain interior bounds for the gradient of solutions. In order to find these bounds we will assume one of the following two conditions on  $g$ :

- (a) There exists  $t_0 > 0$  such that  $g$  is differentiable for  $t \geq t_0$  and

$$\frac{g'(t)}{g(t)^2} \leq C t^{-\gamma}, \quad t \geq t_0$$

for some  $\gamma > 2$ , or

- (b) There exists a positive constant  $C$  such that  $|g(t)| \leq Ct^2$ , for large  $t$ .

Note that condition (a) is verified by the case  $g(s) = s^2 \log(s + 1)$  while (b) is the standard condition to get this kind of interior bounds.

Before proceeding to state our principal results below, it is important to stress that in all of them we will assume the following general hypotheses on  $f$  and  $g$ :

$$(f_0 - g_0) \quad \begin{array}{l} f \text{ and } g \text{ are continuous increasing functions} \\ \text{such that } f(0) = g(0) = 0. \end{array}$$

A word of caution: we are always concerned with nonnegative solutions, but since  $f$  and  $g$  are only continuous, the strict positivity of the solutions cannot be guaranteed, that is, the strong maximum principle is not always valid (cf. [22] and [21] for its validity in this context). In our present situation the solutions could have a “dead core” in the interior of the domain, but we will not be concerned with this aspect of the problem.

We start considering problem  $(P_+)$ , and introduce

$$(1.5) \quad \Gamma(s) = \int_0^{2s} g(t) dt + 2Ns^2.$$

Then we have the following existence/nonexistence result.

**Theorem 1.** *Let  $f$  and  $g$  be functions satisfying  $(f_0 - g_0)$ . Then:*

(i) *If  $f$  does not verify the Keller-Osserman condition (1.2) or if*

$$(1.6) \quad \int_1^\infty \frac{ds}{g^{-1}(f(s))} = \infty,$$

*then problem  $(P_+)$  does not admit nonnegative solutions.*

(ii) *If  $g$  verifies one of the conditions (a) or (b) above and*

$$(1.7) \quad \int_1^\infty \frac{ds}{\Gamma^{-1}(\frac{1}{2}F(s))} < \infty,$$

*where  $\Gamma$  is given by (1.5), then there exists at least a nonnegative solution  $u$  to  $(P_+)$ , which in addition verifies  $u \in C^{1,\alpha}(\Omega)$  for every  $\alpha \in (0, 1)$ .*

Let us now turn to problem  $(P_-)$ . We have already mentioned that the power  $q = 2$  is critical among power-like nonlinearities regarding existence. In the general setting, we will prove that the “criticality” of  $g$  for existence of solutions with large boundary condition depends on the divergence of the integral  $\int_1^\infty \frac{s}{g(s)} ds$ , that is

$$(1.8) \quad \int_1^\infty \frac{s}{g(s)} ds = \infty.$$

**Theorem 2.** *Let  $f$  and  $g$  be functions satisfying  $(f_0 - g_0)$ , and assume that  $g$  verifies condition (1.8). Then:*

(i) *If*

$$(1.9) \quad \int_1^\infty \frac{ds}{f(s) + g(s)} = \infty,$$

*then problem  $(P_-)$  does not have nonnegative solutions.*

(ii) *If  $f$  verifies the Keller-Osserman condition (1.2) or  $f$  verifies  $\lim_{t \rightarrow \infty} f(t) = \infty$  and  $g$  verifies one of the conditions (a) or (b) above and is such that*

$$(1.10) \quad \int_1^\infty \frac{ds}{g(s)} < \infty.$$

then problem  $(P_-)$  admits at least a nonnegative solution  $u$ , which verifies  $u \in C^{1,\alpha}(\Omega)$  for every  $\alpha \in (0, 1)$ .

*Remark 1.* Some relations between the different conditions appeared above are expected to hold. For instance, it is easy to check that (1.9) implies (1.8), while (1.8) is also implied by (b). On the other hand, (a) with the additional condition  $\lim_{t \rightarrow \infty} g(t) = \infty$  gives (1.10).

Our proofs of existence and nonexistence for nonnegative solutions to  $(P_{\pm})$  rely in comparison with solutions to the same problems in balls of  $\mathbb{R}^N$ . Thus it is important to analyze the radial version of those problems, which is in turn close to the one-dimensional version. Since these equations are not integrable, our method inspired in [21] will be to compare them with some integrable ones and this leads to the previous integral conditions of Keller-Osserman type involving  $f$  and  $g$ .

In this work we also are able to prove that condition (1.8) is sharp in the sense that non existence holds not only for  $(P_-)$  but also for the problem

$$(1.11) \quad \begin{cases} \Delta u - g(|\nabla u|) = f(u) & \text{in } \Omega, \\ u = n & \text{on } \partial\Omega, \end{cases}$$

when  $n$  is large enough, provided that the integral in (1.8) converges, see Theorem 6 in Section 5.

Also we find some *uniqueness* results for nonnegative solutions to  $(P_+)$  and  $(P_-)$  under some extra assumptions on  $f$  and  $g$ , which are used to obtain the precise asymptotic behavior of all solutions near the boundary, and are essentially the usual hypotheses in the literature when  $g = 0$  (cf. for instance [7]). See Theorems 8 and 9, respectively, in Section 6. All our results are illustrated by means of examples. Moreover, we complete the study in [4] of the particular choice  $f(s) = s^p$  and  $g(s) = s^q$ , proving uniqueness when one has existence and finding a new range for the exponent  $p$  and  $q$  where non existence holds, see Corollary 13 in Section 7.

The rest of the paper is organized as follows: in Section 2 we deal with some preliminary properties of radial solutions to the Cauchy problems related to  $(P_{\pm})$ . Section 3 is devoted to obtain existence of solutions to the finite boundary value problems associated to  $(P_{\pm})$  while in Section 4 the existence and nonexistence results are considered. In Section 5 we show that condition (1.8) is necessary for existence in problems  $(P_-)$  and (1.11). The uniqueness issue is undertaken in Section 6 while in Section 7 some illustrative examples are analyzed. Finally we include an Appendix where we prove some interior gradient bounds for solutions to  $(P_{\pm})$ .

## 2. PROPERTIES OF RADIAL SOLUTIONS

In this section we are going to prove some preliminary properties of solutions to the Cauchy problem

$$(2.1) \quad \begin{cases} u'' + \frac{N-1}{r}u' = f(u) \pm g(u'), \\ u(0) = u_0, \quad u'(0) = 0, \end{cases}$$

where  $f$  and  $g$  are functions satisfying  $(f_0 - g_0)$ , and  $u_0 \geq 0$ . By continuity, it is well known that there exists at least a solution to (2.1). We are interested only in nonnegative solutions.

Our main result in this section is the following:

**Proposition 3.** *Let  $u$  be a nonnegative nontrivial solution to (2.1). If  $u_0 > 0$  then  $u'(r) > 0$ ,  $u''(r) \geq 0$  for  $r > 0$ . If  $u_0 = 0$ , then there exists  $r_0 \geq 0$  such that  $u \equiv 0$  in  $[0, r_0]$ ,  $u'(r) > 0$ ,  $u''(r) \geq 0$  for  $r > r_0$ . In particular,  $u$  and  $u'$  are nondecreasing functions and for every  $R > 0$  such that  $u$  is defined in  $(0, R)$ ,*

$$(2.2) \quad u(r) \leq u_0 + Ru'(r), \quad r \in (0, R).$$

*Remark 2.* Note that when  $f$  and  $g$  are locally Lipschitz, standard ode's theory implies  $r_0 = 0$ .

*Proof.* Assume first  $u_0 > 0$ . From the equation we obtain  $u''(0) = \frac{1}{N}f(u_0) > 0$ , so that  $u''(r) > 0$  if  $r > 0$ ,  $r$  close enough to zero. This implies  $u'(r) > 0$  for  $r > 0$  close enough to zero. Assume there exists  $r_1 > 0$  with  $u'(r) > 0$  if  $r \in (0, r_1)$  and  $u'(r_1) = 0$ . Then we would have  $u''(r_1) \leq 0$  and the equation would give

$$u''(r_1) = f(u(r_1)) > 0,$$

a contradiction. Hence  $u'(r) > 0$  for  $r > 0$ .

Suppose next  $u_0 = 0$ . Define

$$r_0 = \sup\{\tilde{r} : u(r) = 0 \text{ in } [0, \tilde{r}]\},$$

and let us prove that  $u'(r) > 0$  when  $r > r_0$ . Notice first that there exists a sequence  $r_n \downarrow r_0$  such that  $u'(r_n) > 0$ . If not, we would have  $u' \leq 0$  in  $[r_0, r_0 + \varepsilon]$  for some  $\varepsilon > 0$  and this leads to  $u = 0$  in  $[r_0, r_0 + \varepsilon]$ , contradicting the definition of  $r_0$ .

We may assume  $u(r_n) > 0$ , since in the case  $u(r_n) = 0$  we could take  $\bar{r}_n > r_n$ , close to  $r_n$  with  $u(\bar{r}_n) > 0$ . Now a similar reasoning as before shows that  $u'(r) > 0$  if  $r > r_n$ . Hence  $u'(r) > 0$  for  $r > r_0$ .

Let us now deal with the sign of  $u''$ . We need different proofs for problem (2.1) with a plus or with a minus sign. Let us begin with the minus sign:

$$u'' + \frac{N-1}{r}u' = f(u) - g(u').$$

Assume  $u''(r_2) < 0$  for some  $r_2 > 0$ . Let

$$\tilde{r}_0 = \inf\{\tilde{r} : u''(r) < 0 \text{ in } (\tilde{r}, r_2)\}.$$

Since  $u''(0) \geq 0$ , we have  $u''(\tilde{r}_0) = 0$ . Moreover,  $u''(r) < 0$  for  $r > \tilde{r}_0$ ,  $r$  close to  $\tilde{r}_0$ . This implies that  $u'$  is decreasing for  $r \geq \tilde{r}_0$ ,  $r$  close to  $\tilde{r}_0$ . Since  $u$  is increasing, we have that

$$u'' = f(u) - \frac{N-1}{r}u' - g(u')$$

is increasing. But then  $u''(\tilde{r}_0) = 0$  implies that  $u'' > 0$  if  $r > \tilde{r}_0$ ,  $r$  close to  $\tilde{r}_0$ , a contradiction. Thus  $u'' \geq 0$ .

Next, let us deal with the plus sign:

$$u'' + \frac{N-1}{r}u' = f(u) + g(u').$$

For every  $r > r_0$ , there exists  $r_3 \in (r_0, r)$  such that  $u''(r_3) > 0$ . If not, we would have  $u'' \leq 0$  in  $(r_0, r)$  and since  $u'(r_0) = 0$ , we arrive at  $u' \leq 0$  in  $(r_0, r)$ , which is impossible. Let us assume that  $r_4 = \inf\{s > r_3 : u''(s) = 0\}$  exists. Then, since  $u'' + \frac{N-1}{r}u'$  is increasing in  $(r_3, r_4)$ , we have for sufficiently small  $h > 0$ :

$$(N-1) \left( \frac{u'(r_4)}{r_4} - \frac{u'(r_4-h)}{r_4-h} \right) \geq u''(r_4-h) - u''(r_4) > 0.$$

Dividing by  $h$  and letting  $h \rightarrow 0$  we obtain

$$-\frac{u'(r_4)}{r_4^2} \geq 0,$$

which is impossible. Hence  $u''(s) > 0$  if  $s > r_3$ , and we are done since  $r_0 < r_3 < r$  and  $r > r_0$  is arbitrary.

Finally, since  $u'' \geq 0$ , we have that  $u'$  is nondecreasing. Thus, for  $r \in (0, R)$ :

$$u(r) = u_0 + \int_0^r u'(s) ds \leq u_0 + Ru'(r).$$

This concludes the proof.  $\square$

### 3. EXISTENCE OF SOLUTIONS WITH FINITE DATUM

In this section we assume that  $f$  and  $g$  are functions satisfying  $(f_0 - g_0)$ . Here we will prove that, under suitable conditions on the growth of  $g$ , there always exists a solution to the problems with finite datum

$$(P_{\pm}^n) \quad \begin{cases} \Delta u \pm g(|\nabla u|) = f(u) & \text{in } \Omega, \\ u = n & \text{on } \partial\Omega, \end{cases}$$

where  $n \in \mathbb{N}$ . We warn the reader that  $g$  may have a superquadratic growth in the gradient, thus to ensure existence of solutions we will truncate the nonlinearity, and then look for right bounds for the gradient, in the spirit of [40] (see also [4]).

Since in what follows it will be easier to work with  $|\nabla u|^2$  instead of  $|\nabla u|$ , we denote  $\tilde{g}(t) = g(\sqrt{t})$  for  $t \geq 0$ . Now choose  $M > 0$  and let  $g_M$  be an increasing, bounded function such that  $g_M(t) = \tilde{g}(t)$  for  $t \in [0, M]$ . Consider the truncated problems

$$(3.1) \quad \begin{cases} \Delta u \pm g_M(|\nabla u|^2) = f(u) & \text{in } \Omega, \\ u = n & \text{on } \partial\Omega. \end{cases}$$

Since  $\underline{u} = 0$  is a subsolution to (3.1) while  $\bar{u} = n$  is a supersolution, the result in [16] ensures the existence of a solution  $u \in H^1(\Omega)$  which verifies  $0 \leq u \leq n$ , and then  $0 \leq f(u) \leq f(n)$ . In particular,  $u \in L^\infty(\Omega)$ , and by standard regularity theory  $u \in C^{1,\alpha}(\bar{\Omega})$  (Corollary 8.35 in [30]). According to the comparison principle (see for instance Lemma 2.1 (ii) in [22]), this solution is unique.

In order to prove that the solution  $u$  to (3.1) so obtained solves  $(P_{\pm}^n)$ , we need to show that a suitable value of  $M$  can be selected. This will be accomplished by means of uniform bounds for the gradient. Indeed, if we consider  $\hat{f}(t) = f(t)$  if  $t \in [0, n]$  and  $\hat{f}(t) = f(n)$  if  $t > n$ , it suffices with obtaining bounds only on  $\partial\Omega$ , as the following lemma shows.

**Lemma 4.** *Let  $u$  be the solution to (3.1). Then*

$$|\nabla u| \leq \sup_{\partial\Omega} |\nabla u|.$$

*Proof.* The proof relies on an application of the maximum principle to an equation satisfied by  $|\nabla u|^2$ . But we first regularize the problem: let  $\{f_k\}_{k=1}^\infty, \{g_k\}_{k=1}^\infty \subseteq C^\infty(\mathbb{R})$  such that  $f_k$  and  $g_k$  are increasing for all  $k$  and  $f_k \rightarrow \hat{f}$ ,  $g_k \rightarrow g_M$  uniformly on compact sets of  $\mathbb{R}$ . We may moreover assume that the functions  $f_k, g_k$  are uniformly bounded. Consider the problems

$$(3.2) \quad \begin{cases} \Delta u \pm g_k(|\nabla u|^2) = f_k(u) & \text{in } \Omega, \\ u = n & \text{on } \partial\Omega. \end{cases}$$

Arguing as before, there exists a unique solution  $u_k \in C^{1,\alpha}(\overline{\Omega})$  to (3.2). Indeed, it follows by classical regularity that  $u \in C^2(\overline{\Omega}) \cap C^\infty(\Omega)$ . Let  $v_k = |\nabla u_k|^2$ . A calculation shows that

$$\Delta v_k = 2 \sum_{i,j=1}^N (\partial_{ij} u_k)^2 + 2 \sum_{i=1}^N \partial_i u_k \Delta(\partial_i u_k).$$

On the other hand, differentiating in (3.2) with respect to  $x_i$ , we obtain

$$\Delta(\partial_i u_k) \pm g'_k(|\nabla u_k|^2) \partial_i v_k = f'_k(u_k) \partial_i u_k,$$

so that

$$\Delta v_k \pm 2g'_k(v_k) \nabla u_k \nabla v_k = 2 \sum_{i,j=1}^N (\partial_{ij} u_k)^2 + 2f'_k(u_k) v_k \geq 0 \quad \text{in } \Omega.$$

Thanks to the maximum principle, we have  $v_k \leq \sup_{\partial\Omega} v_k$ , that is,

$$(3.3) \quad |\nabla u_k|^2 \leq \sup_{\partial\Omega} |\nabla u_k|^2.$$

Our next purpose is passing to the limit in (3.3). Since  $f_k, g_k$  are uniformly bounded, we obtain that  $\Delta u_k$  is uniformly bounded in  $\overline{\Omega}$ , and thanks to Theorem 8.33 in [30],

$$|u_k|_{C^{1,\alpha}(\overline{\Omega})} \leq C$$

for some positive constant  $C$ , not depending on  $k$ , and some fixed  $\alpha \in (0, 1)$ . Passing to a subsequence, we have  $u_k \rightarrow u_0$  in  $C^1(\overline{\Omega})$ , and therefore  $u_0$  is a solution to (3.1). By uniqueness,  $u_0 = u$ , and in particular  $u_k \rightarrow u$  in  $C^1(\overline{\Omega})$ . We may pass to the limit in (3.3) to get the lemma proved.  $\square$

To obtain bounds for the gradient of  $u$  we need thus to estimate it on  $\partial\Omega$ . We will achieve this by constructing a suitable subsolution. Notice that, thanks to the regularity of  $\Omega$ , it verifies a uniform exterior sphere condition, and hence there exists  $R_1 > 0$  such that, for every  $x_0 \in \partial\Omega$  we can find  $z_0 \notin \overline{\Omega}$  with  $\overline{B}(z_0, R_1) \cap \partial\Omega = \{x_0\}$ . For  $R_2 > R_1$  let  $A$  be the annulus  $\{x \in \mathbb{R}^N : R_1 < |x - z_0| < R_2\}$ .

Set  $D = A \cap \Omega$  and denote  $\Gamma_1 = \{x \in \mathbb{R}^N : |x - z_0| = R_1\}$ ,  $\Gamma_2 = \{x \in \mathbb{R}^N : |x - z_0| = R_2\}$ . Assume that the problem

$$(3.4) \quad \begin{cases} \Delta v \pm g(|\nabla v|) = f(v) & \text{in } A, \\ v = n & \text{on } \Gamma_1, \\ v = 0 & \text{on } \Gamma_2, \end{cases}$$

admits a radial subsolution  $v \in C^1(\overline{A})$  and choose  $M > \sup_{R_1 \leq r \leq R_2} v'(r)^2$ . In that case,  $g(|\nabla v|) = g_M(|\nabla v|^2)$ , and since  $v \leq u$  on  $\partial D$ , we have by comparison  $v \leq u$  in  $D$ . Moreover,  $v(x_0) = n = u(x_0)$ , so that

$$\frac{\partial u}{\partial \nu}(x_0) \leq \frac{\partial v}{\partial \nu}(x_0).$$

Now notice that  $u = n$  on  $\partial\Omega$  implies, by Hopf's principle, that  $\frac{\partial u}{\partial \nu}(x_0) > 0$ , and thus  $\partial\Omega$  is a level set for  $u$ . This entails that  $\nu$  is parallel to  $\nabla u$  and hence  $|\nabla u(x_0)| = \frac{\partial u}{\partial \nu}(x_0)$ . It follows that

$$|\nabla u(x_0)|^2 \leq v'(R_1)^2 < M.$$

In conclusion,  $u$  is a solution to  $(P_{\pm}^n)$ , as we wanted to see.

Thus the important point in the proof of existence of nonnegative solutions to  $(P_{\pm}^n)$  is the obtention of a radial subsolution to (3.4), that is, a function  $v \in C^1([R_1, R_2])$  verifying

$$(3.5)^{\pm} \quad \begin{cases} v'' + \frac{N-1}{r} v' \pm g(|v'|) \geq f(v) & \text{in } R_1 < r < R_2, \\ v(R_1) = n, \\ v(R_2) = 0, \end{cases}$$

in the weak sense. For this aim, the cases with a plus sign and with a minus sign in  $(3.5)^{\pm}$  have to be analyzed separately. It turns out that the problem with the + sign is substantially simpler, since no further growth conditions on  $f$  nor  $g$  need to be imposed. On the other hand, the obtention of a subsolution for problem (3.5) with a minus sign is much more involved, and it strongly depends on the function  $g$ . In this case, we need to assume condition (1.8) since we will show in Section 5 that it is necessary for existence of solutions either to  $(P_-)$  or to  $(P_-^n)$  when  $n$  is large enough.

**Proposition 5.** *Let  $f, g$  be functions satisfying  $(f_0 - g_0)$ . Then*

- i)  $(P_+^n)$  admits a unique nonnegative solution  $u_n \in C^{1,\alpha}(\bar{\Omega})$  for every  $n \in \mathbb{N}$ .*
- ii) If (1.8) holds,  $(P_-^n)$  admits a unique nonnegative solution  $u_n$  for every  $n \in \mathbb{N}$ .*

Moreover, in both cases one has that  $0 < u_n < n$  in  $\Omega$  and  $u_n \in C^{1,\alpha}(\bar{\Omega})$  for every  $\alpha \in (0, 1)$ .

*Proof.* We start proving *i)*. To obtain a subsolution to  $(3.5)^+$ , it suffices with choosing the unique nonnegative solution to

$$\begin{cases} \Delta v = f(v) & \text{in } A, \\ v = n & \text{on } \Gamma_1, \\ v = 0 & \text{on } \Gamma_2, \end{cases}$$

which is easily constructed by means of the method of sub and supersolutions, by taking  $\underline{v} = 0$ ,  $\bar{v} = n$ . Thus the existence of a solution to  $(P_+^n)$  is obtained thanks to the previous discussion.

Now we prove *ii)*. As before, it suffices with constructing a radial function verifying  $(3.5)^-$ . Thus we need a function  $v \in C^1([R_1, R_2])$  verifying

$$(3.6) \quad \begin{cases} (r^{N-1} v')' \geq r^{N-1} (f(v) + g(|v'|)), \\ v(R_1) = n, \\ v(R_2) = 0. \end{cases}$$

With the change of variables

$$s = \begin{cases} \log r & \text{if } N = 2, \\ -\frac{1}{N-2} \frac{1}{r^{N-2}} & \text{if } N \geq 3, \end{cases}$$

and letting  $w(s) = v(r)$ , the inequality (3.6) gets transformed into:

$$(3.7) \quad \begin{cases} w'' \geq h(s) \left( f(w) + g\left(\frac{1}{r^{N-1}} |w'|\right) \right), \\ w(a) = n, \\ w(b) = 0, \end{cases}$$

where  $a = \log R_1$ ,  $b = \log R_2$  if  $N = 2$ , while  $a = -\frac{1}{N-2} \frac{1}{R_1^{N-2}}$ ,  $b = -\frac{1}{N-2} \frac{1}{R_2^{N-2}}$  when  $N \geq 3$ . The function  $h(s) = r^{2(N-1)}$  and  $'$  stands now for differentiation with respect to

s. To obtain (3.7), it is enough to have

$$\begin{cases} w'' \geq R_2^{2(N-1)} \left( f(w) + g \left( R_1^{-(N-1)} |w'| \right) \right), \\ w(a) = n, \\ w(b) = 0. \end{cases}$$

Our intention is keeping  $R_1$  fixed (recall that it comes from the uniform exterior sphere condition) and treating  $R_2$  as a parameter. Thus if we choose  $\bar{R} > R_1$  and take  $R_2 \leq \bar{R}$ , we are looking for a solution to

$$(3.8) \quad \begin{cases} w'' = \bar{R}^{2(N-1)} \left( f(w) + g \left( R_1^{-(N-1)} |w'| \right) \right), \\ w(a) = n, \\ w(b) = 0. \end{cases}$$

It is easily seen that solutions to (3.8) are decreasing. If we set  $z(s) = w(b - s)$ , then we look for an increasing function  $z$  which solves

$$\begin{cases} z'' = \bar{R}^{2(N-1)} \left( f(z) + g \left( R_1^{-(N-1)} z' \right) \right), \\ z(0) = 0, \\ z(b - a) = n. \end{cases}$$

It is therefore natural to analyze the initial value problem

$$(3.9) \quad \begin{cases} z'' = \bar{R}^{2(N-1)} \left( f(z) + g \left( R_1^{-(N-1)} z' \right) \right), \\ z(0) = 0, \\ z'(0) = z_0, \end{cases}$$

for  $z_0 > 0$  and see if it is possible to choose  $z_0$  so that  $z(b - a) = n$ .

For  $z_0 > 0$ , problem (3.9) admits a minimal solution, which is defined in an interval  $[0, T)$  with  $T \leq \infty$ . Moreover, when  $T < \infty$  we have  $z(s) \rightarrow \infty$  or  $z'(s) \rightarrow \infty$  as  $s \rightarrow T^-$ . Notice that solutions are increasing and convex. They are also increasing with  $z_0$ .

We claim that condition (1.8) implies that  $z(s) \rightarrow \infty$  as  $s \rightarrow T$  (that is, the solution can not cease to exist because of the blow-up of the derivative). If this were not the case, we would have  $z(s) \rightarrow \bar{z}$  as  $s \rightarrow T$  for a certain finite  $\bar{z}$  and  $z'(s) \rightarrow \infty$  as  $s \rightarrow T$ . Let us rule out this possibility. If we multiply the equation by  $z'$ , taking into account that  $z$  is bounded we have

$$\frac{z' z''}{A + B g \left( R_1^{-(N-1)} z' \right)} \leq z',$$

for some positive constants  $A$  and  $B$ . Integrating in  $[0, s)$  for  $s$  close to  $T$  and then letting  $s \rightarrow T$ , we obtain, with a change of variables in the integral:

$$\int_{R_1^{-(N-1)} z_0}^{\infty} \frac{\tau}{A + B g(\tau)} d\tau \leq R_1^{-(N-1)} \bar{z},$$

which contradicts condition (1.8). Hence, we have shown that  $T < \infty$  implies  $z(s) \rightarrow \infty$  (and thus also  $z'(s) \rightarrow \infty$ ) as  $s \rightarrow T$ .

Denote by  $T(z_0)$  the maximal interval of existence of the minimal solution to (3.9). Two options may occur: either  $T(z_0) = \infty$  for every  $z_0 > 0$ , that is, the minimal solution is always global, or there exists  $z_1 > 0$  such that  $T(z_1) < \infty$ . In the first case, choosing  $\delta > 0$  small enough, and since  $z(s) \geq z_0 s$  by convexity, we have  $z(\delta) \geq z_0 \delta > n$ , provided  $z_0$  is large enough. Hence there exists  $s_0 \in (0, \delta)$  such that  $z(s_0) = n$ . We may now choose

$R_2$  such that  $b - a = s_0$ , and the existence of a function verifying (3.5) is shown in this case.

In the remaining case  $T(z_1) < \infty$  for some  $z_1 > 0$ , we have, since solutions are increasing with  $z_0$ , that  $T(z_0) < \infty$  for every  $z_0 > z_1$ . In particular, for every  $z_0 > z_1$  there exists  $s_0 \in (0, T(z_0))$  such that  $z(s_0) = n$ . Since  $z(s_0) \geq z_0 s_0$ , we also have  $s_0 \rightarrow 0$  as  $z_0 \rightarrow \infty$ . We choose as before a large value of  $z_0$  and then  $R_2$  so that  $b - a = s_0$ , and in this way we have constructed the desired subsolution. This concludes the proof.  $\square$

#### 4. EXISTENCE AND NONEXISTENCE OF SOLUTIONS TO $(P_{\pm})$

This section is dedicated to prove Theorems 1 and 2, that is, existence and nonexistence results for problems  $(P_{\pm})$ . Let us first comment on the method of proof. To show the existence of a solution to either problem, we consider the solutions  $u_n$  to the finite problems  $(P_{\pm}^n)$  furnished by Proposition 5. If  $B \subset \Omega$  is an arbitrary ball, then we obtain by the comparison principle in Lemma 2.1 (ii) of [22] that  $u_n \leq u_{n,B}$ , where  $u_{n,B}$  is the unique solution to  $(P_{\pm}^n)$  in the ball  $B$ . Assume we prove

$$(4.1) \quad \sup_{n \in \mathbb{N}} u_{n,B}(x) < \infty \quad x \in B.$$

This would imply that  $u_n$  is locally uniformly bounded in  $\Omega$ , and since the sequence  $u_n$  is increasing we have  $u_n \rightarrow u := \sup_{n \in \mathbb{N}} u_n$  point-wise in  $\Omega$ . Thanks to Theorem A.1 in the Appendix, we would have that  $|\nabla u_n|$  is locally uniformly bounded. Then we would obtain that  $\Delta u_n = h_n$ , for a function  $h_n$  which is locally uniformly bounded. Using classical regularity (for instance (4.45) in [30]), we obtain bounds for  $u_n$  in  $C^{1,\alpha}(\Omega)$  for every  $\alpha \in (0, 1)$ . It is then standard to conclude by means of a diagonal procedure that, passing to a subsequence,  $u_n \rightarrow u$  in  $C^1(\Omega)$ . Then  $u$  is a solution to  $(P_{\pm})$ , and by standard regularity  $u \in C^{1,\alpha}(\Omega)$  for every  $\alpha \in (0, 1)$ . We remark that in the previous reasoning the radius of the ball  $B$  can be taken as small as desired.

As for the nonexistence issue, let  $B$  be a large ball containing  $\Omega$  and denote again by  $u_{n,B}$  the solution to  $(P_{\pm}^n)$  in  $B$ . If either problem  $(P_+)$  or  $(P_-)$  had a solution, we would obtain by comparison that  $u \geq u_{n,B}$ . We would arrive at a contradiction if we prove that

$$(4.2) \quad u_{n,B} \rightarrow \infty \quad \text{uniformly in } B.$$

Thus it is clear that only the radial case needs to be dealt with.

With these ideas in mind, we proceed to prove Theorems 1 and 2.

*Proof of Theorem 1.* We need only prove (4.1) or (4.2). For notational simplicity we will drop the subindex  $B$  and denote the solution to  $(P_{\pm}^n)$  in the ball  $B$  by  $u_n$ . Notice that  $u_n$  has to be radially symmetric, and thus it verifies

$$(4.3) \quad \begin{cases} u'' + \frac{N-1}{r} u' = f(u) - g(|u'|), \\ u'(0) = 0, \\ u(R) = n, \end{cases}$$

where  $'$  stands for derivative with respect to  $r = |x|$ . Denote by  $u_{0,n} = u_n(0) = \min u_n$ .

Let us prove part (i), that is, nonexistence of solutions for  $(P_+)$ . Assume first  $f$  does not verify the Keller-Osserman condition (1.2). Since, according to Proposition 3,  $u' > 0$

we have from (4.3) that  $u'' \leq f(u)$ , so that multiplying by  $u'$  and integrating in  $(0, r)$  we arrive at  $u' \leq \sqrt{2F(u)}$ , where  $F$  is the primitive of  $f$  vanishing at zero. Thus

$$\int_{u_{0,n}}^n \frac{ds}{\sqrt{2F(s)}} \leq R.$$

Letting  $n \rightarrow \infty$  we obtain that  $u_{0,n} \rightarrow \infty$  since  $f$  does not verify (1.2). Thus (4.2) holds and this shows nonexistence.

Next, suppose condition (1.6) holds. Thanks to Proposition 3, we have  $u', u'' \geq 0$ , so that (4.3) implies  $f(u) \geq g(u')$ . Hence

$$\frac{u'}{g^{-1}(f(u))} \leq 1,$$

and integrating in  $(r_0, R)$  for some  $r_0$  close to  $R$ :

$$\int_{u_n(r_0)}^n \frac{ds}{g^{-1}(f(s))} \leq R.$$

Letting  $n \rightarrow \infty$  and using condition (1.6) we deduce again that  $u_n(r_0) \rightarrow \infty$ , so that  $u_{0,n} \rightarrow \infty$ , and hence no nonnegative solutions to  $(P_+)$  exist.

Let us prove now the existence result in part (ii). We first show that condition (1.7) allows us to construct a radial supersolution in a ball  $B$  which blows up on  $\partial B$ , provided that the radius  $R$  of the ball is small enough. We will assume for the moment that  $R \leq 1/2$ , and will search for a supersolution in the form

$$\bar{u}(r) = \phi(R^2 - r^2),$$

where  $\phi(0) = \infty$ . It is not hard to show that  $\bar{u}$  will be a supersolution if

$$4r^2\phi'' - 2N\phi' + g(2r|\phi'|) \leq f(\phi),$$

where  $'$  stands for differentiation with respect to  $t = R^2 - r^2$ . Assume for the moment that  $\phi' < 0$ ,  $\phi'' > 0$ . Then it suffices to have

$$(4.4) \quad \phi'' - 2N\phi' + g(-\phi') \leq f(\phi).$$

Let us now choose the function  $\phi$ . Thanks to condition (1.7), the problem

$$(4.5) \quad \begin{cases} \Gamma(|\phi'|) = \frac{1}{2}F(\phi) & t > 0, \\ \phi(0) = \infty, \end{cases}$$

admits a unique solution. It is more or less standard to check that  $\phi' < 0$  (and moreover  $\phi(t), \phi'(t) \rightarrow 0$  as  $t \rightarrow \infty$ ). Notice that the convergence of the integral in (1.7) implies

$$\frac{\Gamma^{-1}\left(\frac{1}{2}F(s)\right)}{s} \rightarrow \infty \quad (s \rightarrow \infty),$$

and hence

$$\frac{|\phi'(t)|}{|\phi(t)|} \rightarrow \infty \quad (t \rightarrow 0).$$

In particular, there exists  $\varepsilon > 0$  such that  $\phi(t) \leq |\phi'(t)|$  if  $0 < t \leq \varepsilon$ . Restrict  $R$  further to have  $R^2 \leq \varepsilon$ , so that  $t = R^2 - r^2 \leq \varepsilon$ . Let us see that  $\phi$  verifies the required properties. Taking derivative in (4.5) we have

$$\phi'' = \frac{1}{2}f(\phi) \frac{-\phi'}{2g(-2\phi') + 4N(-\phi')}.$$

We deduce then that  $\phi'' > 0$  and

$$(4.6) \quad \phi'' \leq \frac{1}{2} f(\phi).$$

On the other hand, we have by the monotonicity of  $f$  and  $g$ :

$$F(t) = \int_0^t f(s) ds \leq f(t)t$$

and

$$\Gamma(t) = \int_0^{2t} g(s) ds + 2Nt^2 \geq \int_t^{2t} g(s) ds + 2Nt^2 \geq tg(t) + 2Nt^2.$$

Then

$$(4.7) \quad g(-\phi') - 2N\phi' \leq \frac{\Gamma(-\phi')}{-\phi'} = \frac{1}{2} \frac{F(\phi)}{-\phi'} \leq \frac{1}{2} \frac{f(\phi)\phi}{-\phi'} \leq \frac{1}{2} f(\phi).$$

Adding (4.6) and (4.7) we obtain (4.4).

To summarize, we have constructed a supersolution  $\bar{u}$  to (4.3) in  $B$ , with  $\bar{u} = \infty$  on  $\partial B$ , with the only restriction that  $R$  is small enough. It follows by comparison that  $u_{n,B} \leq \bar{u}$  in  $B$ , and hence (4.1) follows, provided only that  $R$  is sufficiently small. Thus existence of a nonnegative solution to  $(P_+)$  is proved in this case.  $\square$

*Proof of Theorem 2.* To show part (i), we prove (4.2). Recall that, the solution  $u_{n,B}$  to  $(P_{\pm}^n)$  verifies the Cauchy problem

$$(4.8) \quad \begin{cases} u'' + \frac{N-1}{r} u' = f(u) + g(|u'|), \\ u'(0) = 0, \\ u(0) = u_{0,n}, \end{cases}$$

and thanks to Proposition 3, it verifies  $u'(r) \geq 0$ ,  $u(r) \leq u_0 + Ru'(r)$ . Hence if we assume that  $R \geq 1$ , we also have  $u'' \leq f(u) + g(u') \leq f(u_{0,n} + Ru') + g(u') \leq f(u_{0,n} + Ru') + g(u_{0,n} + Ru')$ . It follows after an integration in  $(r_0, R)$  for some  $r_0$  close to  $R$  that

$$\int_{u'(r_0)}^{u'(R)} \frac{ds}{f(u_{0,n} + Rs) + g(u_{0,n} + Rs)} \leq R.$$

Let us set  $u_{0,n} + Rs = \tau$ ; since the solution to  $(P_{\pm}^n)$  verifies  $n = u(R) \leq u_{0,n} + Ru'(R)$ , we arrive at

$$\int_{u_{0,n}}^n \frac{d\tau}{f(\tau) + g(\tau)} \leq 1.$$

We mention in passing that  $u_{0,n} = 0$  is only possible when the integral converges at zero, so no problem arises in this case (cf. a related situation in Theorem 1.2 in [21]). The divergence of the integral implies  $u_{0,n} \rightarrow \infty$  as  $n \rightarrow \infty$ , and (4.2) gets proved. Thus no solutions exist under condition (1.9).

To show the existence result in part (ii), we argue as in Theorem 1 and prove (4.1). Notice that, if  $f$  verifies the Keller-Osserman condition (1.2), and since  $u_{n,B}$  satisfies  $\Delta u \geq f(u)$  in  $B$ , we obtain by comparison that  $u_{n,B} \leq U$ , where  $U$  is the minimal solution to  $\Delta U = f(U)$  in  $B$ , with  $U = \infty$  on  $\partial B$ . Thus (4.1) is immediate in this case.

So assume  $\lim_{t \rightarrow \infty} f(t) = \infty$  and  $g$  verifies condition (1.10). Since the equation in (4.8) can be written as  $(r^{N-1} u')' = r^{N-1}(f(u) + g(u'))$ , we can integrate in  $(0, r)$  for an arbitrary  $r$  and use that  $u$  and  $u'$  are increasing (Proposition 3) to obtain

$$\begin{aligned} u'(r) &= \frac{1}{r^{N-1}} \int_0^r s^{N-1} (f(u(s)) + g(u'(s))) ds \\ &\leq \frac{r}{N} (f(u(r)) + g(u'(r))). \end{aligned}$$

Taking this inequality to (4.8) we obtain that

$$(4.9) \quad u'' \geq \frac{1}{N} (f(u) + g(u')) \geq \frac{1}{N} (f(u_{0,n}) + g(u')).$$

When  $u_{0,n} > 0$ , it follows by integrating in  $(0, R)$  and performing the standard change of variables  $s = u'(r)$  in the integral that

$$(4.10) \quad \frac{1}{N} R \leq \int_0^{u'(R)} \frac{1}{f(u_{0,n}) + g(s)} ds \leq \int_0^\infty \frac{1}{f(u_{0,n}) + g(s)} ds.$$

If we had  $u_{0,n} \rightarrow \infty$  the last integral in (4.10) would tend to zero by dominated convergence and we would reach a contradiction. Thus  $u_{0,n}$  remains bounded and this shows (4.1), as was to be proved.  $\square$

### 5. NECESSITY OF CONDITION (1.8)

In this section we will prove that condition (1.8) is necessary for problem  $(P_-)$  to have a solution. Moreover, it is also necessary for problem  $(P_-^n)$  when  $n$  is sufficiently large. This result is in the spirit of the nonexistence example in [51] (see Theorem 1 in Chapter III, §16 there).

Thus our main result here is:

**Theorem 6.** *Let  $f$  and  $g$  be functions satisfying  $(f_0 - g_0)$ , and assume that  $\lim_{t \rightarrow \infty} f(t) = \infty$ . If condition (1.8) does not hold, that is,*

$$(5.1) \quad \int_1^\infty \frac{s}{g(s)} ds < \infty,$$

*then problem  $(P_-)$  does not have any nonnegative solution. Moreover, there exists  $n_0 = n_0(\Omega)$  such that the problem with finite boundary datum  $(P_-^n)$  does not have nonnegative solutions  $u \in C^1(\bar{\Omega})$  for  $n \geq n_0$ .*

*Remark 3.* A slight modification of the proof of Theorem 6 shows that if  $h \in C^1(\partial\Omega)$  is positive, the problem

$$\begin{cases} \Delta u - g(|\nabla u|) = f(u) & \text{in } \Omega, \\ u = h & \text{on } \partial\Omega, \end{cases}$$

has no nonnegative solutions  $u \in C^1(\bar{\Omega})$  if  $|h|_\infty$  is large enough.

Let us begin by considering the particular instance where  $\Omega = B_R$ . In this case, we will show that no radial solutions exist in both situations.

**Lemma 7.** *Assume  $g$  verifies (5.1). Then problem  $(P_-)$  does not have radial nonnegative solutions. Moreover, the problem with finite datum*

$$(5.2) \quad \begin{cases} \Delta u - g(|\nabla u|) = f(u) & \text{in } B_R, \\ u = n & \text{on } \partial B_R, \end{cases}$$

*does not have nonnegative solutions if  $n$  is large.*

*Proof.* Let us show first that  $(P_-)$  does not admit radial nonnegative solutions. It follows from (4.9) in the proof of Theorem 2 that

$$u'u'' \geq \frac{1}{N}(f(u) + g(u'))u' \geq \frac{1}{N}g(u')u'.$$

Thus if we divide by  $g(u')$  and integrate between  $r_0$  and  $r$  for some arbitrary  $r_0 \in (0, R)$  and  $r$  close to  $R$ , we obtain

$$\int_{u'(r_0)}^{u'(r)} \frac{s}{g(s)} ds \geq \frac{1}{N}(u(r) - u(r_0)),$$

and we arrive at a contradiction when we let  $r \rightarrow R$  thanks to (5.1). Hence no radial nonnegative solutions to  $(P_-)$  can exist.

Let us turn now to the proof that no solutions to (5.2) exist when  $n$  is large enough. We assume that there exists a sequence  $n_k \rightarrow \infty$  such that each problem (5.2) with  $n = n_k$  has a nonnegative solution  $u_k$ . By uniqueness, this solution must be radial, and hence it verifies

$$\begin{cases} u'' + \frac{N-1}{r}u' = f(u) + g(|u'|), \\ u'(0) = 0, \\ u(R) = n_k. \end{cases}$$

Let  $u_{0,k} = u_k(0)$ . Since we may assume that  $n_k$  is increasing in  $k$  and solutions to the associated initial value problem are also increasing with respect to the initial datum, we have that  $u_{0,k}$  is increasing. If  $u_{0,k} \rightarrow \infty$ , we would obtain (4.10), leading to a contradiction as in the proof of Theorem 2. Thus  $u_{0,k}$  is bounded.

Let  $r_k = \inf\{r \in (0, R) : u'_k(r) > 1\}$ . Observe that  $r_k$  exists for large  $k$ , for otherwise  $u'_k \leq 1$  in  $(0, R)$ , and this yields the contradiction  $n_k \leq R + u_{0,k}$ . It follows that  $u_k(r_k)$  is bounded, so that, arguing as in the first part of the proof:

$$\int_1^{u'_k(R)} \frac{s}{g(s)} ds \geq \frac{1}{N}(n_k - u_k(r_k)),$$

leading to a contradiction with (5.1). Thus there can be no solutions to (5.2) when  $n$  is large enough, and this concludes the proof.  $\square$

Let us now prove Theorem 6. The proof relies in showing that if a solution to either  $(P_-)$  or to  $(P_-^n)$  exists in a smooth bounded domain  $\Omega$ , then we can construct a solution for the same problem in a suitable ball  $B_R$ , contradicting Lemma 7. The argument to do this is similar to the one used in Section 3: we truncate the function  $g$ , find a solution to the truncated problem and then show that it is a solution to the original problem by obtaining bounds for the gradient.

*Proof of Theorem 6.* Assume first that for some  $n$  there exists a solution  $u \in C^1(\overline{\Omega})$  to

$$(5.3) \quad \begin{cases} \Delta u - g(|\nabla u|) = f(u) & \text{in } \Omega, \\ u = n & \text{on } \partial\Omega. \end{cases}$$

Take an arbitrary ball  $B_R \subset \Omega$  which is tangent to  $\partial\Omega$  at some  $x_0 \in \partial\Omega$ . For  $M > \sup_{\overline{\Omega}} |\nabla u|^2$ , consider the truncated problem

$$\begin{cases} \Delta v - g_M(|\nabla v|^2) = f(v) & \text{in } B_R, \\ v = n & \text{on } \partial B_R, \end{cases}$$

where  $g_M$  is a bounded function with  $g_M(t^2) = g(t)$  if  $t^2 \leq M$ . As in Section 3, there exists a nonnegative solution  $v$  to this problem by means of the method of sub and supersolutions. By uniqueness this solution is radial and by Proposition 3, it verifies  $v' \geq 0$ , while  $v'$  is nondecreasing.

On the other hand, we have by comparison that  $u < v$  in  $B_R$ , and since  $u(x_0) = v(x_0)$ :

$$\frac{\partial v}{\partial \nu}(x_0) \leq \frac{\partial u}{\partial \nu}(x_0),$$

so that

$$v'(R)^2 \leq |\nabla u(x_0)|^2 < M.$$

Since  $v'$  is increasing and positive we have  $v'(r)^2 < M$  for  $0 \leq r \leq R$ , so that  $v$  is a solution to (5.2), contradicting Lemma 7 if  $n$  is large enough (depending only on  $R$ ).

Finally, let us tackle the question of nonexistence for problem  $(P_-)$ . Assume there exists a solution  $u$  to  $(P_-)$ . Take a ball  $B_R(x_0) \subset \Omega$  tangent to  $\partial\Omega$  at some point  $z_0$ . For small  $\varepsilon > 0$  let  $x_\varepsilon = x_0 - \varepsilon\nu(z_0)$ , where  $\nu(z_0)$  is the outward unit normal to  $\partial\Omega$  at  $z_0$ , so that  $B_R(x_\varepsilon) \subset\subset \Omega$ . Denote  $n = n(\varepsilon) = \sup_{\partial B_R(x_\varepsilon)} u$ . Then

$$\begin{cases} \Delta u - g(|\nabla u|) = f(u) & \text{in } B_R(x_\varepsilon), \\ u \leq n & \text{on } \partial B_R(x_\varepsilon), \end{cases}$$

and an argument like in the first part of the proof shows that the problem

$$\begin{cases} \Delta v - g(|\nabla v|) = f(v) & \text{in } B_R(x_\varepsilon), \\ v = n & \text{on } \partial B_R(x_\varepsilon), \end{cases}$$

has a solution. However, when  $\varepsilon \rightarrow 0$ ,  $n \rightarrow \infty$ , and we arrive at a contradiction with the already proved nonexistence of solutions to (5.3) for large  $n$ . This concludes the proof.  $\square$

## 6. UNIQUENESS

In this section we consider some results concerning uniqueness of nonnegative solutions to  $(P_\pm)$ . As we mentioned earlier, uniqueness is achieved thanks to some special monotonicity of the nonlinearities  $f$  and  $g$ , together with the knowledge of the exact boundary behavior of all possible nonnegative solutions. To obtain this boundary behavior, we will assume that  $f$  verifies the Keller-Osserman condition (1.2), and another condition which is usual in the literature, namely:

$$(6.1) \quad \limsup_{t \rightarrow \infty} \frac{\psi_f(\lambda t)}{\psi_f(t)} < 1$$

for every  $\lambda > 1$ , where

$$(6.2) \quad \psi_f(t) = \int_t^\infty \frac{ds}{\sqrt{F(s)}}$$

(cf. [7]). Let us mention that this condition holds for instance when  $f(t)/t^p$  is increasing for some  $p$  and large  $t$ , as can be easily checked.

We prove the following theorems:

**Theorem 8.** *Let  $f$  and  $g$  be functions satisfying  $(f_0 - g_0)$  and assume that  $f$  verifies (1.2), (6.1),  $f(t)/t$  is increasing and  $g(t)/t$  is decreasing. Then problem  $(P_+)$  admits a unique nonnegative solution.*

**Theorem 9.** *Let  $f$  and  $g$  be functions satisfying  $(f_0 - g_0)$ . Assume that  $g$  verifies (1.8) and (6.1) with  $\psi_f$  replaced by  $\psi_g$ , where*

$$(6.3) \quad \psi_g(t) = \int_t^\infty \frac{ds}{g(s)}.$$

*Assume moreover that  $f(t)/t$  and  $g(t)/t$  are increasing and that*

$$\lim_{t \rightarrow \infty} \frac{f(t)}{g(t)} = 0.$$

*Then problem  $(P_-)$  has a unique nonnegative solution.*

Let us begin with problem  $(P_+)$ . The boundary behavior is given in the next lemma, where we denote  $d(x) = \text{dist}(x, \partial\Omega)$ .

**Lemma 10.** *Assume  $f$  and  $g$  verify  $(f_0 - g_0)$  and that  $f$  verifies (1.2). Assume moreover that  $g$  verifies*

$$(6.4) \quad \limsup_{t \rightarrow \infty} \frac{g(t)}{t} < \infty.$$

*Then every nonnegative solution  $u$  to  $(P_+)$  is such that*

$$(6.5) \quad \lim_{d(x) \rightarrow 0} \frac{\psi_f(u(x))}{d(x)} = 1,$$

*where  $\psi_f$  is given by (6.2). If in addition  $f$  verifies (6.1), then*

$$(6.6) \quad \lim_{d(x) \rightarrow 0} \frac{u(x)}{\phi_f(d(x))} = 1,$$

*where  $\phi_f$  is the inverse function of  $\psi_f$ .*

*Proof.* The proof of boundary estimates relies on comparison with solutions in balls and annuli. For this sake, we need to analyze first the radial case. Thus let  $u$  be a solution to

$$\begin{cases} u'' + \frac{N-1}{r} u' = f(u) - g(|u'|), \\ u'(0) = 0, \\ u(R) = \infty. \end{cases}$$

It follows from Proposition 3 that  $u'' \geq 0$ ,  $u' > 0$ , so that  $u'' \leq f(u)$ . Multiplying by  $u'$  and integrating we obtain  $u' \leq \sqrt{2F(u)}$ , so that in a standard way:

$$\int_{u(r)}^\infty \frac{ds}{\sqrt{2F(s)}} \leq R - r,$$

that is,

$$\limsup_{r \rightarrow R} \frac{\psi_f(u(r))}{R - r} \leq 1.$$

To obtain the complementary inequality, we first observe that since  $f$  verifies the Keller-Osserman condition, it follows that

$$\lim_{t \rightarrow \infty} \frac{\sqrt{F(t)}}{f(t)} = 0$$

(see the Appendix in [6]) and thus condition (6.4) implies that for some positive constant  $C$ :

$$\frac{g(u'(r))}{f(u(r))} \leq C \frac{u'(r)}{f(u(r))} \leq C \frac{\sqrt{2F(u(r))}}{f(u(r))} \rightarrow 0 \quad \text{as } r \rightarrow R.$$

Now notice that  $\frac{u'(r)}{r} \leq \frac{\sqrt{2F(u)}}{r_0}$  if  $r \geq r_0$ , where  $r_0 \in (0, R)$  is arbitrary. Thus

$$\begin{aligned} u'' &\geq f(u) - g(u') - \frac{N-1}{r_0} \sqrt{2F(u)} \\ &\geq f(u) \left( 1 - \frac{g(u')}{f(u)} - \frac{N-1}{r_0} \frac{\sqrt{2F(u)}}{f(u)} \right) \\ &\geq (1 - \varepsilon) f(u) \end{aligned}$$

for some small  $\varepsilon$ , provided  $r$  is close enough to  $R$ . An integration as before provides

$$\liminf_{r \rightarrow R} \frac{\psi_f(u(r))}{R-r} \geq 1.$$

Thus (6.5) holds in this case. It is shown with only minor modifications that it also holds when the domain  $\Omega$  is an annulus.

Now let  $u$  be a solution to  $(P_+)$  in a smooth bounded domain  $\Omega$ . Choose a radius  $R > 0$  such that  $\partial\Omega$  verifies the uniform interior sphere condition. For  $x \in \Omega$  with  $d(x) < R$ , let  $\bar{x} \in \partial\Omega$  be its projection onto  $\partial\Omega$ . There exists  $z_{\bar{x}} \in \Omega$  such that the ball  $B_R(z_{\bar{x}})$  is contained in  $\Omega$  and is tangent to  $\partial\Omega$  at  $\bar{x}$ . Let  $u_B$  be a solution to  $(P_+)$  in this ball. We obtain by comparison:

$$u(x) \leq u_B(|x - z_{\bar{x}}|),$$

for  $x \in B_R(z_{\bar{x}})$ . Using  $d(x) = |x - \bar{x}| = R - |x - z_{\bar{x}}|$ , and since  $\psi_f$  is decreasing,

$$(6.7) \quad \frac{\psi_f(u(x))}{d(x)} \geq \frac{\psi_f(u_B(|x - z_{\bar{x}}|))}{d(x)} = \frac{\psi_f(u_B(|x - z_{\bar{x}}|))}{R - |x - z_{\bar{x}}|}.$$

Thanks to (6.5) in the radial case, we know that for a small  $\varepsilon > 0$  there exists  $\delta > 0$  such that for  $d(x) < \delta$  the last term in (6.7) is  $\geq 1 - \varepsilon$ . Hence

$$\frac{\psi_f(u(x))}{d(x)} \geq 1 - \varepsilon \quad \text{when } d(x) < \delta.$$

This immediately gives

$$\liminf_{d(x) \rightarrow 0} \frac{\psi_f(u(x))}{d(x)} \geq 1.$$

For the complementary inequality we use the uniform exterior sphere condition. Given  $x \in \Omega$  close to  $\partial\Omega$ , we take its projection  $\bar{x} \in \partial\Omega$ . There exist  $R' > 0$  (not depending on  $x$ ) and  $w_{\bar{x}} \notin \bar{\Omega}$  such that  $B_{R'}(w_{\bar{x}}) \cap \Omega = \emptyset$ ,  $\overline{B_{R'}(w_{\bar{x}})} \cap \bar{\Omega} = \{\bar{x}\}$ . Take  $R'' \gg 1$  so that  $\Omega \subset B_{R''}(w_{\bar{x}})$ .

Then  $\Omega \subseteq A := B_{R''}(w_{\bar{x}}) \setminus \overline{B_{R'}(w_{\bar{x}})}$  and taking any nonnegative solution  $u_A$  to  $(P_+)$  in the annulus  $A$  we obtain by comparison  $u(x) \geq u_A(|x - w_{\bar{x}}|)$ , so that

$$\frac{\psi_f(u(x))}{d(x)} \leq \frac{\psi_f(u_A(|x - w_{\bar{x}}|))}{|x - w_{\bar{x}}| - R'} \leq 1 + \varepsilon,$$

when  $d(x) < \delta$ . Thus (6.5) is proved.

Finally, when condition (6.1) holds, it is well known that (6.5) implies (6.6) (see [7]).  $\square$

We now prove the uniqueness result for problem  $(P_+)$ . The proof is an adaptation of the argument in [26].

*Proof of Theorem 8.* Let  $u, v$  be arbitrary nonnegative solutions to  $(P_+)$ . Notice that under the assumptions on  $g$  we have (6.4), so that Lemma 10 can be applied and it gives

$$\lim_{d(x) \rightarrow 0} \frac{u(x)}{v(x)} = 1.$$

Choose  $\varepsilon > 0$ . Then there exists  $\delta > 0$  so that

$$(6.8) \quad (1 - \varepsilon)v(x) \leq u(x) \leq (1 + \varepsilon)v(x)$$

for  $d(x) \leq \delta$ . Let  $\Omega^\delta = \{x \in \Omega : d(x) > \delta\}$  and consider the problem

$$(6.9) \quad \begin{cases} \Delta w + g(|\nabla w|) = f(w) & \text{in } \Omega^\delta, \\ w = u & \text{on } \partial\Omega^\delta, \end{cases}$$

which has as its unique solution  $w = u$ . Now the monotonicity of  $f(t)/t$  and  $g(t)/t$  implies that  $(1 + \varepsilon)v$  is a supersolution to (6.9), while  $(1 - \varepsilon)v$  is a subsolution. It follows by comparison that  $(1 - \varepsilon)v \leq u \leq (1 + \varepsilon)v$  in  $\Omega^\delta$ . Hence the inequality (6.8) holds throughout  $\Omega$ , and we can let  $\varepsilon \rightarrow 0$  to obtain  $u = v$ . This shows uniqueness.  $\square$

Now let us turn once again to problem  $(P_-)$ . We have an analogue of Lemma 10:

**Lemma 11.** *Assume  $f$  and  $g$  verify  $(f_0 - g_0)$ ,  $g$  verifies (1.8) and (6.1) holds with  $\psi_f$  replaced by  $\psi_g$ , where  $\psi_g$  is given by (6.3). If  $f$  satisfies in addition*

$$(6.10) \quad \lim_{t \rightarrow \infty} \frac{f(t)}{g(t)} = 0,$$

then every nonnegative solution to  $(P_-)$  verifies

$$(6.11) \quad u(x) \sim \int_{d(x)}^1 \phi_g(t) dt$$

as  $d(x) \rightarrow 0$ , where  $\phi_g$  is the inverse function of  $\psi_g$ .

*Proof.* The proof of (6.11) is obtained by comparison with solutions in balls and annuli, as in Lemma 10. Thus we only prove it in the case  $\Omega = B_R$ . If  $u$  is a radial nonnegative solution, then

$$(6.12) \quad u'' + \frac{N-1}{r}u' = f(u) + g(u') \geq g(u').$$

Notice that  $u''$  is not necessarily increasing, but the group  $u'' + \frac{N-1}{r}u'$  is, thanks to Proposition 3. Thus for every  $r_0 \in (0, R)$  and  $r \in (r_0, R)$ :

$$\begin{aligned} u'(r) &= u'(r_0) + \int_{r_0}^r u''(s) ds \leq u'(r_0) + \int_{r_0}^r \left( u''(s) + \frac{N-1}{s}u'(s) \right) ds \\ &\leq u'(r_0) + (r - r_0) \left( u''(r) + \frac{N-1}{r}u'(r) \right) \\ &\leq u'(r_0) + (N-1) \frac{R-r_0}{r_0} u'(r) + (R-r_0)u''(r). \end{aligned}$$

Taking  $r_0$  close enough to  $R$  so that  $(N-1) \frac{R-r_0}{r_0} \leq \frac{1}{2}$ , we obtain

$$u'(r) \leq 2u'(r_0) + 2(R-r_0)u''(r).$$

Dividing by  $u''(r)$ , letting  $r \rightarrow R$  and then  $r_0 \rightarrow R$ , we obtain  $u'(r)/u''(r) \rightarrow 0$  as  $r \rightarrow R$ . Thus from (6.12):

$$(1 + \varepsilon)u'' \geq g(u'),$$

when  $r$  is close enough to  $R$ . Dividing by  $g(u')$  and integrating in  $(r, R)$  for  $r$  close to  $R$ :

$$\int_{u'(r)}^{\infty} \frac{ds}{g(s)} \geq \frac{1}{1 + \varepsilon}(R - r).$$

Hence we arrive at

$$\liminf_{r \rightarrow R} \frac{\psi_g(u'(r))}{R - r} \geq 1.$$

On the other hand, since  $u'$  is increasing, it is shown as before that  $u(r)/u'(r) \rightarrow 0$  as  $r \rightarrow R$ , and in particular  $u \leq u'$  if  $r$  is close to  $R$ . Thanks to condition (6.10) we also have  $f(t) \leq \varepsilon g(t)$  when  $t$  is large enough. Then

$$u'' \leq u'' + \frac{N-1}{r} u' = f(u) + g(u') \leq f(u') + g(u') \leq (1 + \varepsilon)g(u'),$$

and it follows that

$$\int_{u'(r)}^{\infty} \frac{ds}{g(s)} \leq (1 + \varepsilon)(R - r).$$

We then obtain

$$\lim_{r \rightarrow R} \frac{\psi_g(u'(r))}{R - r} = 1.$$

With the additional condition (6.1) with  $\psi_g$  we also have

$$\lim_{r \rightarrow R} \frac{u'(r)}{\phi_g(R - r)} = 1,$$

as in Lemma 10. Thanks to l'Hôpital rule we deduce that

$$u(r) \sim u(0) + \int_0^r \phi_g(R - s)ds = u(0) + \int_{R-r}^R \phi_g(t)dt \sim \int_{R-r}^1 \phi_g(t)dt,$$

which is (6.11) in a ball of radius  $R$ . This concludes the proof.  $\square$

*Remark 4.* We notice that  $\phi'_g = -g(\phi_g)$ , so that

$$\int_{R-r}^1 \phi_g(t)dt = - \int_{R-r}^1 \frac{\phi_g(t)\phi'_g(t)}{g(\phi_g(t))} dt = \int_{\phi_g(1)}^{\phi_g(R-r)} \frac{s}{g(s)} ds,$$

and the boundary behavior (6.11) can be written as

$$u(x) \sim \int_1^{\phi_g(d(x))} \frac{s}{g(s)} ds$$

when  $d(x) \rightarrow 0$ .

We close this section by mentioning that the proof of Theorem 9 is a slight variation of that of Theorem 8, using Lemma 11 instead of Lemma 10, and therefore it will not be given.

## 7. SOME EXAMPLES

In this section we quote some important cases of nonlinearities  $f$  and  $g$ , and particularize the results of our paper to them. It is easily seen that conditions (a) or (b) in the Introduction hold for all of them. Let us begin with  $(P_+)$ .

*7.1.  $(P_+)$  with  $g(t) = t^q$  for some  $q > 0$ .* In this case  $g^{-1}(t) = t^{1/q}$ , while  $\Gamma(s) = \frac{(2s)^{q+1}}{q+1} + 2Ns^2$ . Thus when  $0 < q \leq 1$  we have  $\Gamma(s) \sim \text{constant} \cdot s^2$  as  $s \rightarrow \infty$ , so that  $\Gamma^{-1}(s) \sim \text{constant} \cdot \sqrt{s}$  as  $s \rightarrow \infty$ , and condition (1.7) is nothing more than Keller-Osserman condition (1.2). When  $q > 1$ , on the contrary,  $\Gamma(s) \sim \text{constant} \cdot s^{q+1}$  as  $s \rightarrow \infty$ , so that  $\Gamma^{-1}(s) \sim \text{constant} \cdot s^{\frac{1}{q+1}}$  and (1.7) reads as

$$\int_1^\infty \frac{ds}{F(s)^{\frac{1}{q+1}}} < \infty.$$

So that we obtain directly from Theorem 1:

**Corollary 12.** *Let  $f$  be an increasing continuous function with  $f(0) = 0$ . Then for  $0 < q \leq 1$ , the problem*

$$(7.1) \quad \begin{cases} \Delta u + |\nabla u|^q = f(u) & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

*admits a nonnegative solution if and only if  $f$  verifies (1.2), while for  $q > 1$ , (7.1) admits a nonnegative solution when*

$$(7.2) \quad \int_1^\infty \frac{ds}{F(s)^{\frac{1}{q+1}}} < \infty$$

*and does not have any nonnegative solution when*

$$(7.3) \quad \int_1^\infty \frac{ds}{f(s)^{\frac{1}{q}}} = \infty.$$

Let us mention that conditions (7.2) and (7.3) are not exhaustive. This is easily seen by taking  $f(t) = t^q(\log t)^\alpha$ , for  $\alpha$  verifying  $q < \alpha \leq q + 1$ , where neither condition (7.2) nor (7.3) hold.

As particular cases of Corollary 12, let us single out the functions  $f(t) = t^p$ ,  $p > 0$  and  $f(t) = e^t - 1$ , where all possibilities are exhausted. The next two corollaries complement the results in [4], since uniqueness and nonexistence were not considered there.

**Corollary 13.** *Let  $p, q > 0$ . For the problem*

$$\begin{cases} \Delta u + |\nabla u|^q = u^p & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

*we have:*

- (i) *If  $0 < q \leq 1$ , there exists a nonnegative solution if and only if  $p > 1$ , and it is unique.*
- (ii) *If  $q > 1$ , there exists a nonnegative solution if and only if  $p > q$ .*

**Corollary 14.** *The problem*

$$\begin{cases} \Delta u + |\nabla u|^q = e^u - 1 & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

*admits a nonnegative solution for every  $q > 0$ . This solution is unique when  $0 < q \leq 1$ .*

7.2. ( $P_+$ ) with  $g(t) = e^t - 1$ . This is an interesting example since we are analyzing a problem with a huge growth in the gradient. Notice that  $\Gamma(s) \sim e^{2s}$  for large  $s$  so that  $\Gamma^{-1}(s) \sim \frac{1}{2} \log s$  as  $s \rightarrow \infty$ . Since also  $g^{-1}(t) \sim \log t$  we immediately have from Theorem 1 that the condition

$$(7.4) \quad \int_1^\infty \frac{ds}{\log F(s)} < \infty$$

implies existence, while

$$(7.5) \quad \int_1^\infty \frac{ds}{\log f(s)} = \infty$$

gives nonexistence. The important remark is that conditions (7.4) and (7.5) are complementary. This is easily seen by noticing that  $F(2t) \geq f(t)$  for  $t \geq 1$ , so that the divergence of the integral in (7.4) implies (7.5) and the convergence of the integral in (7.5) implies condition (7.4). Thus:

**Corollary 15.** *Let  $f$  be a continuous increasing function verifying  $f(0) = 0$ . Then the problem*

$$\begin{cases} \Delta u + e^{|\nabla u|} - 1 = f(u) & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

*admits a nonnegative solution if and only if*

$$\int_1^\infty \frac{ds}{\log f(s)} < \infty.$$

Let us finally consider problem ( $P_-$ ).

7.3. ( $P_-$ ) with  $g(t) = t^q$  for some  $q > 0$ . Condition (1.8) is equivalent to  $q \leq 2$ , while (1.10) in Theorem 2 means  $q > 1$ . On the other hand, condition (1.9), which in our present situation translates into

$$(7.6) \quad \int_1^\infty \frac{ds}{f(s) + s^q} = \infty,$$

is equivalent, when  $0 < q \leq 1$ , to

$$(7.7) \quad \int_1^\infty \frac{ds}{f(s)} = \infty.$$

Indeed, it is clear that (7.6) implies (7.7). For the other implication, if we assume that (7.6) does not hold, since  $f(s) + s^q$  is increasing, we have that  $\lim_{s \rightarrow \infty} \frac{s}{f(s) + s^q} = 0$ , which implies that  $\lim_{s \rightarrow \infty} \frac{s}{f(s)} = 0$ , so that  $f(s) \sim f(s) + s^q$  as  $s \rightarrow \infty$ , and then (7.7) does not hold. Hence

**Corollary 16.** *Let  $f$  be increasing and continuous. Assume moreover that  $f(0) = 0$  and  $\lim_{t \rightarrow \infty} f(t) = \infty$ . Then for  $1 < q \leq 2$  there exists at least a nonnegative solution to*

$$(7.8) \quad \begin{cases} \Delta u - |\nabla u|^q = f(u) & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega. \end{cases}$$

*When  $0 < q \leq 1$  and  $f$  verifies (1.2) there exists a nonnegative solution to (7.8), and there exists none when*

$$\int_1^\infty \frac{ds}{f(s)} = \infty.$$

As before, the two important cases  $f(t) = t^p$ ,  $p > 0$  and  $f(t) = e^t - 1$  give:

**Corollary 17.** *Let  $p, q > 0$ . Then the problem*

$$\begin{cases} \Delta u - |\nabla u|^q = u^p & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

*has no nonnegative solutions if either  $q > 2$  or if  $p, q \leq 1$ . When  $1 < q \leq 2$  or  $p > 1$ , there exists a nonnegative solution. Moreover, the solution is unique if  $1 \leq p < q \leq 2$ .*

**Corollary 18.** *Let  $q > 0$ . The problem*

$$\begin{cases} \Delta u - |\nabla u|^q = e^u - 1 & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

*has no nonnegative solution if  $q > 2$ , while it has at least one when  $0 < q \leq 2$ .*

*Remark 5.* As in the case considered in Section 7.1, the conditions in Corollary 16 are not exhaustive. For instance, the corollary cannot be applied to the function  $f(t) = t(\log t)^\alpha$  when  $1 < \alpha \leq 2$ .

*7.4. ( $P_-$ ) with  $g(t) = t^2 \log(t+1)$ .* This is in our opinion one of the most interesting cases in our study, since it consists in an equation with a superquadratic growth in the gradient. It is not hard to see that condition (1.8) is verified. Moreover, also condition (1.10) in Theorem 2 holds, so that:

**Corollary 19.** *Assume  $f$  is an increasing continuous function verifying  $f(0) = 0$  and  $\lim_{t \rightarrow \infty} f(t) = \infty$ . Then the problem*

$$\begin{cases} \Delta u - |\nabla u|^2 \log(|\nabla u| + 1) = f(u) & \text{in } \Omega, \\ u = \infty & \text{on } \partial\Omega, \end{cases}$$

*admits a nonnegative solution. If moreover  $\frac{f(t)}{t}$  is increasing and*

$$\lim_{t \rightarrow \infty} \frac{f(t)}{t^2 \log t} = 0,$$

*then the solution is unique.*

## APPENDIX: GRADIENT BOUNDS

It is important for the existence proofs in Section 4 to dispose of interior gradient bounds for solutions to the equations

$$(A.1) \quad \Delta u \pm g(|\nabla u|) = f(u) \quad \text{in } \Omega.$$

These bounds can be obtained independently of the growth of the function  $g$ . However, we need to impose restrictions (a) or (b) in the introduction, which are of technical nature.

By modifying the arguments in the Appendix to [35], where the case  $g(t) = t^q$ ,  $q > 0$  was considered, we are going to prove:

**Theorem A. 1.** *Let  $f$  and  $g$  be increasing continuous functions with  $f(0) = g(0) = 0$ . Assume  $g$  verifies one of the following two conditions:*

(a) *There exists  $t_0 > 0$  such that  $g$  is differentiable for  $t \geq t_0$  and*

$$\frac{g'(t)}{g(t)^2} \leq C t^{-\gamma}, \quad t \geq t_0$$

*for some  $\gamma > 2$ , or*

(b) *There exists a positive constant  $C$  such that  $|g(t)| \leq Ct^2$ , for large  $t$ .*

Let  $u \in C^1(\Omega)$  be a nonnegative solution to (A.1). Then for every pair of smooth subdomains  $\Omega' \subset\subset \Omega'' \subset\subset \Omega$  there exists a constant  $C$  depending only on  $|f(u)|_{L^\infty(\Omega'')}$ ,  $\Omega'$ ,  $\Omega''$  and  $g$  such that

$$(A.2) \quad \sup_{\Omega'} |\nabla u| \leq C.$$

*Proof.* As in the proof of Lemma 4, since it is easier to deal with  $|\nabla u|^2$  than with  $|\nabla u|$ , we set  $g(t) = \tilde{g}(t^2)$ .

Let us begin with case (b), which is more or less classical. If  $u \in C^1(\Omega)$  is a solution to (A.1), we have  $\Delta u \in L^\infty_{\text{loc}}(\Omega)$ , so that by standard regularity  $u \in W_{\text{loc}}^{2,p}(\Omega)$  for every  $p > 1$ . We may use Theorem 6.5 in Chapter IV of [33] to obtain that for  $\Omega' \subset\subset \Omega$ , there exists  $\alpha \in (0, 1)$  and a constant  $M > 0$  which only depends on  $|u|_{L^\infty(\Omega')}$ ,  $|f(u)|_{L^\infty(\Omega')}$ , the constant  $C$  in (b) and the distance from  $\Omega'$  to  $\partial\Omega$ , such that

$$|u|_{C^{1,\alpha}(\overline{\Omega}')} \leq M.$$

This shows (A.2) in case (b).

As for case (a), notice that we may assume  $\lim_{t \rightarrow \infty} g(t) = \infty$ , otherwise (b) holds. The condition on  $g$  implies

$$(A.3) \quad \frac{\tilde{g}'(t)}{\tilde{g}(t)^2} \leq C t^{-\frac{1+\gamma}{2}}$$

for large  $t$  and some positive constant  $C$ . By direct integration it is also easy to see that this entails

$$(A.4) \quad \tilde{g}(t) \geq C t^{\frac{\gamma-1}{2}}$$

for large  $t$  and some positive  $C$ .

By approximation, as in Lemma 4, we may assume that  $\tilde{g}$  and  $f$  are  $C^1$ . More precisely, let  $\{f_k\}, \{g_k\}$  be sequences of  $C^1$  functions such that  $f_k \rightarrow f, g_k \rightarrow \tilde{g}$  uniformly in compacts of  $\mathbb{R}$ . We may also assume that  $f_k$  is increasing and  $g_k$  verifies condition (a) with a uniform constant  $C$  (notice that  $g$  is  $C^1$  for large  $t$ , so we could take  $g_k(t) = \tilde{g}(t)$  for large  $t$  if we wish). Let  $\Omega' \subset\subset \Omega'' \subset\subset \Omega$  be smooth subdomains. The problem

$$\begin{cases} \Delta v \pm g_k(|\nabla v|) = f_k(v) & \text{in } \Omega', \\ v = u & \text{on } \partial\Omega', \end{cases}$$

has a unique solution  $v_k \in C^2(\overline{\Omega}')$ . The existence follows by the method of sub and supersolutions and the uniqueness is a consequence of the comparison principle in [22]. Assuming we prove

$$(A.5) \quad \sup_{\Omega'} |\nabla v_k| \leq C$$

for a constant  $C$  which does not depend on  $k$ , we can argue as in Lemma 4 to deduce that (passing to a subsequence)  $v_k \rightarrow u$  in  $C^1(\overline{\Omega}')$ . Letting  $k \rightarrow \infty$  in (A.5) we obtain (A.2) for  $u$ .

Thus we assume  $f$  and  $\tilde{g}$  are  $C^1$ . Next take  $\varphi \in C_0^\infty(\Omega'')$  such that  $\varphi \equiv 1$  in  $\Omega'$ ,  $0 \leq \varphi \leq 1$  and  $|\Delta\varphi| \leq C\varphi^\theta, |\nabla\varphi|^2 \leq C\varphi^{1+\theta}$ , for some  $\theta \in (0, 1)$  to be chosen later.

Let  $w = |\nabla u|^2$ . We have seen in Lemma 4 that  $w$  verifies the equation

$$\Delta w = 2|D^2u|^2 + 2f'(u)w \mp 2\tilde{g}'(w)\nabla u \nabla w$$

in  $\Omega$ , so that the function  $z := \varphi w$  verifies

$$\Delta z = 2\varphi|D^2u|^2 + 2\varphi f'(u)w \mp 2\tilde{g}'(w)\varphi \nabla u \nabla w + 2\nabla w \nabla \varphi + w \Delta \varphi.$$

Now notice that  $z = 0$  outside  $\Omega''$ . Hence there exists  $x_0 \in \Omega''$  such that  $z$  achieves its maximum at  $x_0$ . Since  $\Delta z(x_0) \leq 0$  and  $\nabla z(x_0) = 0$ , we have

$$\begin{aligned}
(A.6) \quad 2\varphi|D^2u|^2 &\leq \mp 2\tilde{g}'(w)\nabla u\nabla\varphi w + \frac{|\nabla\varphi|^2}{\varphi}w - w\Delta\varphi \\
&\leq 2\tilde{g}'(w)|\nabla u||\nabla\varphi|w + \frac{|\nabla\varphi|^2}{\varphi}w + w|\Delta\varphi| \\
&\leq C\tilde{g}'(w)\varphi^{\frac{1+\theta}{2}}w^{\frac{3}{2}} + C\varphi^\theta w,
\end{aligned}$$

at the point  $x_0$ , where we have used  $\tilde{g}' \geq 0$  and  $f' \geq 0$ . On the other hand, thanks to Cauchy-Schwarz inequality, we have

$$(\Delta u)^2 = \left( \sum_{i=1}^N \partial_{ii}u \right)^2 \leq N \sum_{i=1}^N (\partial_{ii}u)^2 \leq N|D^2u|^2,$$

so that (A.6) implies that

$$\frac{2}{N} \varphi(\Delta u)^2 \leq C\tilde{g}'(w)\varphi^{\frac{1+\theta}{2}}w^{\frac{3}{2}} + C\varphi^\theta w$$

at the point  $x_0$ .

Moreover, using the equation (A.1),

$$(\Delta u)^2 = (\tilde{g}(w) \mp f(u))^2 \geq (\tilde{g}(w) - C)^2 \geq C\tilde{g}(w)^2 - C,$$

where  $C$  depends on  $\tilde{g}$  and  $|f(u)|_{L^\infty(\Omega'')}$ , so that

$$\varphi\tilde{g}(w)^2 \leq C\varphi + C\tilde{g}'(w)\varphi^{\frac{1+\theta}{2}}w^{\frac{3}{2}} + C\varphi^\theta w$$

at  $x_0$ . Using (A.3) and  $w = z/\varphi$  we get

$$(A.7) \quad \tilde{g}(w)^2 \leq C + C\tilde{g}(w)^2 z^{1-\frac{\gamma}{2}} \varphi^{\frac{1+\theta}{2}-2+\frac{\gamma}{2}} + C\varphi^{\theta-2}z.$$

We now choose  $\theta \geq 3 - \gamma$ , which is always possible since  $3 - \gamma < 1$ . Then (A.7) becomes

$$(A.8) \quad \tilde{g}(w)^2 \leq C + C\tilde{g}(w)^2 z^{1-\frac{\gamma}{2}} + C\varphi^{\theta-2}z.$$

Now assume  $Cz(x_0)^{1-\frac{\gamma}{2}} \leq 1/2$ , since on the contrary there is nothing to prove. Then (A.8) implies  $\tilde{g}(w)^2 \leq C + C\varphi^{\theta-2}z$  and using (A.4) we arrive at  $w^{\gamma-1} \leq C + C\varphi^{\theta-2}z$ , that is

$$z^{\gamma-1} \leq C\varphi^{\gamma-1} + C\varphi^{\theta+\gamma-3}z \leq C + Cz.$$

Taking into account that  $\gamma > 2$ , we obtain an upper bound for  $z(x_0)$  which only depends on  $|f(u)|_{L^\infty(\Omega'')}$ ,  $\Omega'$ ,  $\Omega''$  and  $g$ . Since  $x_0$  is a point where  $z$  achieves its maximum

$$\varphi|\nabla u|^2 \leq C \quad \text{in } \Omega,$$

and using that  $\varphi \equiv 1$  in  $\Omega'$ , we arrive at (A.2). This concludes the proof.  $\square$

**Acknowledgements.** We would like to thank the referee for a careful reading of the paper and for suggestions which improved its final form. J. G-M was supported by Ministerio de Ciencia e Innovación and FEDER under grant MTM2008-05824 (Spain) and A. Q. was partially supported by Fondecyt Grant # 1070264. All three authors were partially supported by USM Grant # 12.09.17 and Programa Basal CMM, U. de Chile.

## REFERENCES

- [1] A. AFTALION, M. DEL PINO, R. LETELIER, *Multiple boundary blow-up solutions for nonlinear elliptic equations*, Proc. Roy. Soc. Edinburgh **133** (2) (2003), 225–235.
- [2] A. AFTALION, W. REICHEL, *Existence of two boundary blow-up solutions for semilinear elliptic equations*, J. Diff. Eqns. **141**, (1997), 400–421.
- [3] S. ALARCÓN, G. DÍAZ, R. LETELIER, J. M. REY, *Expanding the asymptotic explosive boundary behavior of large solutions to a semilinear elliptic equation*, Nonlinear Anal. **72** (2010), 2426–2443.
- [4] C. BANDLE, E. GIARRUSSO, *Boundary blowup for semilinear elliptic equations with nonlinear gradient terms*, Adv. Differential Equations **1** (1996), 133–150.
- [5] C. BANDLE, M. MARCUS, *'Large' solutions of semilinear elliptic equations: Existence, uniqueness and asymptotic behaviour*, J. Anal. Math. **58** (1992), 9–24.
- [6] C. BANDLE, M. MARCUS, *Asymptotic behaviour of solutions and their derivatives, for semilinear elliptic problems with blowup on the boundary*, Ann. Inst. H. Poincaré Anal. Non Linéaire **12** (1995), no. 2, 155–171.
- [7] C. BANDLE, M. MARCUS, *On second order effects in the boundary behaviour of large solutions of semilinear elliptic problems*, Diff. Int. Eqns. **11** (1998), 23–34.
- [8] C. BANDLE, M. MARCUS, *Dependence of blowup rate of large solutions of semilinear elliptic equations, on the curvature of the boundary*, Complex Var. Theory Appl. **49** (2004), no. 7-9, 555–570.
- [9] L. BIEBERBACH,  *$\Delta u = e^u$  und die automorphen funktionen*, Math. Ann. **77** (1916), 173–212.
- [10] M. CHUAQUI, C. CORTÁZAR, M. ELGUETA, C. FLORES, J. GARCÍA-MELIÁN, R. LETELIER, *On an elliptic problem with boundary blow-up and a singular weight: the radial case*, Proc. Roy. Soc. Edinburgh **133** (2003), 1283–1297.
- [11] M. CHUAQUI, C. CORTÁZAR, M. ELGUETA, J. GARCÍA-MELIÁN, *Uniqueness and boundary behaviour of large solutions to elliptic problems with singular weights*, Comm. Pure Appl. Anal. **3** (2004), 653–662.
- [12] F. CÎRSTEĂ, Y. DU, *General uniqueness results and variation speed for blow-up solutions of elliptic equations*, Proc. London Math. Soc. **91** (2005), 459–482.
- [13] F. CÎRSTEĂ, V. RĂDULESCU, *Uniqueness of the blow-up boundary solution of logistic equations with absorption*, C. R. Acad. Sci. Paris Sér. I Math. **335** (5) (2002), 447–452.
- [14] F. C. CÎRSTEĂ, V. RĂDULESCU, *Asymptotics for the blow-up boundary solution of the logistic equation with absorption*. C. R. Math. Acad. Sci. Paris **336** (2003), no. 3, 231–236.
- [15] M. DEL PINO, R. LETELIER, *The influence of domain geometry in boundary blow-up elliptic problems*, Nonlinear Anal. **48** (6) (2002), 897–904.
- [16] J. DEUEL, P. HESS, *A Criterion for the existence of solutions of non-linear elliptic boundary value problems*, Proc. Roy. Soc. Edinburgh Sect. A **74** (1975), 49–54.
- [17] G. DÍAZ, R. LETELIER, *Explosive solutions of quasilinear elliptic equations: existence and uniqueness*, Nonlinear Anal. **20** (1992), 97–125.
- [18] H. DONG, S. KIM, M. SAFONOV, *On uniqueness of boundary blow-up solutions of a class of nonlinear elliptic equations* Comm. Partial Differential Equations **33** (2008), no. 1-3, 177–188.
- [19] Y. DU, Q. HUANG, *Blow-up solutions for a class of semilinear elliptic and parabolic equations*, SIAM J. Math. Anal. **31** (1999), 1–18.
- [20] S. DUMONT, L. DUPAIGNE, O. GOUBET, V. RĂDULESCU, *Back to the Keller-Osserman condition for boundary blow-up solutions*. Adv. Nonlinear Stud. **7** (2007), no. 2, 271–298.
- [21] P. FELMER, M. MONTENEGRO, A. QUAAS, *A note on the strong maximum principle and the compact support principle*, J. Diff. Eqns. **246** (2009), 39–49.
- [22] P. FELMER, A. QUAAS, *On the strong maximum principle for quasilinear elliptic equations and systems*, Adv. Differential Equations **7** (2002), no. 1, 25–46.
- [23] J. GARCÍA-MELIÁN *Boundary behavior for large solutions to elliptic equations with singular weights*. Nonlinear Anal. **67** (2007), no. 3, 818–826.
- [24] J. GARCÍA-MELIÁN *Uniqueness for boundary blow-up problems with continuous weights*. Proc. Amer. Math. Soc. **135** (2007), no. 9, 2785–2793.
- [25] J. GARCÍA-MELIÁN, *Uniqueness of positive solutions for a boundary blow-up problem*, J. Math. Anal. Appl. **360** (2009), 530–536.
- [26] J. GARCÍA-MELIÁN, R. LETELIER-ALBORNOZ, J. SABINA DE LIS, *Uniqueness and asymptotic behaviour for solutions of semilinear problems with boundary blow-up*, Proc. Amer. Math. Soc. **129** (2001), 3593–3602.

- [27] M. GHERGU, C. NICULESCU, V. RĂDULESCU, *Explosive solutions of elliptic equations with absorption and non-linear gradient term*, Proc. Indian Acad. Sci. Math. Sci. **112** (2002), no. 3, 441–451.
- [28] E. GIARRUSSO, *On blow up solutions of a quasilinear elliptic equation*, Math. Nachr. **213** (2000), 89–104.
- [29] E. GIARRUSSO, *Asymptotic behaviour of large solutions of an elliptic quasilinear equation in a borderline case*, C. R. Acad. Sci. Paris Ser. I Math. **331** (2000), no. 10, 777–782.
- [30] D. GILBARG, N. S. TRUDINGER, “Elliptic Partial Differential Equations of Second Order”. SpringerVerlag, 1983.
- [31] J. B. KELLER, *On solutions of  $\Delta u = f(u)$* , Comm. Pure Appl. Math. **10** (1957), 503–510.
- [32] V. A. KONDRAT’EV, V. A. NIKISHKIN, *Asymptotics, near the boundary, of a solution of a singular boundary value problem for a semilinear elliptic equation*, Differential Equations **26** (1990), 345–348.
- [33] O. A. LADYŽENSKAJA, N. N. URAL’CEVA, “Linear and quasilinear elliptic equations”. Academic Press, New York-London, 1968.
- [34] A. LAIR, A. W. WOOD, *Large solutions of semilinear elliptic equations with nonlinear gradient terms*, Int. J. Mat. & Math. Sci. **22** (1999), 869–883.
- [35] J. M. LASRY, P. L. LIONS, *Nonlinear elliptic equations with singular boundary conditions and stochastic control with state constraints. I. The model problem*, Math. Ann. **283** (1989), 583–630.
- [36] A. C. LAZER, P. J. MCKENNA, *On a problem of Bieberbach and Rademacher*, Nonlinear Anal. **21** (1993), 327–335.
- [37] A. C. LAZER, P. J. MCKENNA, *Asymptotic behaviour of solutions of boundary blow-up problems*, Differential Integral Equations **7** (1994), 1001–1019.
- [38] T. LEONORI, *Large solutions for a class of nonlinear elliptic equations with gradient terms*, Adv. Nonlinear Stud. **7** (2007), no. 2, 237–269.
- [39] T. LEONORI, A. PORRETTA, *The boundary behavior of blow-up solutions related to a stochastic control problem with state constraint*, SIAM J. Math. Anal. **39** (2007/08), no. 4, 1295–1327.
- [40] P. L. LIONS, *Résolution des problèmes elliptiques quasilineaires*, Arch. Rat. Mech. Anal. **74** (1980), 335–353.
- [41] C. LOEWNER, L. NIRENBERG, *Partial differential equations invariant under conformal of projective transformations*, Contributions to Analysis (a collection of papers dedicated to Lipman Bers), Academic Press, New York, 1974, p. 245–272.
- [42] J. LÓPEZ-GÓMEZ, *The boundary blow-up rate of large solutions*, J. Diff. Eqns. **195** (2003), 25–45.
- [43] J. LÓPEZ-GÓMEZ, *Optimal uniqueness theorems and exact blow-up rates of large solutions*, J. Diff. Eqns. **224** (2006), 385–439.
- [44] M. MARCUS, L. VÉRON, *Uniqueness and asymptotic behaviour of solutions with boundary blow-up for a class of nonlinear elliptic equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire **14** (2) (1997), 237–274.
- [45] M. MARCUS, L. VÉRON, *Existence and uniqueness results for large solutions of general nonlinear elliptic equations*, J. Evol. Equ. **3** (2003), no. 4, 637–652.
- [46] R. OSSERMAN, *On the inequality  $\Delta u = f(u)$* , Pac. J. Math. **7** (1957), 1641–1647.
- [47] S. I. POHOZAEV, *The Dirichlet problem for the equation  $\Delta u = u^2$* , Dokl. Akad. Nauk. SSSR **134** (1960), 769–772.
- [48] A. PORRETTA, *Local estimates and large solutions for some elliptic equations with absorption*, Adv. Differential Equations **9** (2004), no. 3-4, 329–351.
- [49] H. RADEMACHER, *Einige besondere Probleme partieller Differentialgleichungen*, in: P. Frank, R. von Mises (Eds.) Die Differential- und Integralgleichungen der Mechanik und Physik I, 2nd. ed., Rosenberg, New York (1943), 838–845.
- [50] V. RĂDULESCU, *Singular phenomena in nonlinear elliptic problems: from boundary blow-up solutions to equations with singular nonlinearities*, in “Handbook of Differential Equations: Stationary Partial Differential Equations”, Vol. **4** (Michel Chipot, Editor) (2007), 483–591.
- [51] J. SERRIN, *The problem of Dirichlet for quasilinear elliptic differential equations with many independent variables*, Philos. Trans. Roy. Soc. London Ser. A **264** (1969), 413–496.
- [52] Z. ZHANG, *Boundary blow-up elliptic problems with nonlinear gradient terms*, J. Diff. Eqns. **228** (2006), no. 2, 661–684.
- [53] Z. ZHANG, *Boundary blow-up elliptic problems of Bieberbach and Rademacher type with nonlinear gradient terms*, Nonlinear Anal. **67** (2007), no. 3, 727–734.
- [54] Z. ZHANG, *Boundary blow-up elliptic problems with nonlinear gradient terms and singular weights*, Proc. Roy. Soc. Edinburgh Sect. A **138** (2008), no. 6, 1403–1424.

S. ALARCÓN AND A. QUAAS

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDAD TÉCNICA FEDERICO SANTA MARÍA  
CASILLA V-110, AVDA. ESPAÑA, 1680 – VALPARAÍSO, CHILE.

*E-mail address:* `salomon.alarcon@usm.cl`, `alexander.quaas@usm.cl`

J. GARCÍA-MELIÁN

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE LA LAGUNA.

C/. ASTROFÍSICO FRANCISCO SÁNCHEZ s/N, 38271 – LA LAGUNA, SPAIN

and

INSTITUTO UNIVERSITARIO DE ESTUDIOS AVANZADOS (IUDEA) EN FÍSICA ATÓMICA,  
MOLECULAR Y FOTÓNICA, FACULTAD DE FÍSICA, UNIVERSIDAD DE LA LAGUNA

C/. ASTROFÍSICO FRANCISCO SÁNCHEZ s/N, 38203 – LA LAGUNA , SPAIN

*E-mail address:* `jjgarmel@ull.es`